

Anticancer Activity and Mechanism of Alpha-Helical Antimicrobial Peptide

Yibing Huang*

Key Laboratory for Molecular Enzymology and Engineering of the Ministry of Education, College of Life Science, Jilin University, China

*Corresponding author: Yibing Huang, Associate Professor at Key Laboratory for Molecular Enzymology and Engineering of the Ministry of Education, College of Life Science, Jilin University, China, Tel: 86-431-8515-5245; E-mail: huangyibing@jlu.edu.cn

Received date: November 22, 2016; Accepted date: November 30, 2016; Published date: December 07, 2016

Copyright: © 2016 Huang Y. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Huang Y (2016) Anticancer Activity and Mechanism of Alpha-Helical Antimicrobial Peptide. Drug Des 5: e133. doi: 10.4172/2169-0138.1000e133

Editorial

Anticancer peptides (ACPs) have become promising molecules as new anticancer agents due to the unique mechanism and several extraordinary properties, such as smaller size, high activity, low immunogenicity, good biocompatibility, etc. [1].

HPRP-A1 is an 15 amino acid residues antimicrobial peptide (AMP), derived from the N-terminus of ribosomal protein L1 (RpL1) of *Helicobacter pylori*. It can be induced to form an amphipathic α -helical structure in hydrophobic medium and exhibited good antimicrobial and antifungal activity [2,3]. Based on the mechanism of antimicrobial peptides, they can describe the integrity of cell membrane by the electrostatic interactions and hydrophobic interaction. HPRP-A1 has been also found that can induce HeLa cell apoptosis by both the extrinsic and intrinsic pathways of the caspase cascade [4]. In order to improve the penetrability and specificity against cancer cell lines, a new hybrid peptide of HPRP-A1-TAT has been designed and synthesized by linked a cell-permeating peptide TAT to the C-terminus of HPRP-A1 [5]. Compared to HPRP-A1, the hybrid peptide of HPRP-A1-TAT exhibited higher anticancer activity against HeLa cells with lower toxicity against human RBC by increasing early apoptosis of HeLa cells and inducing caspase activity. Furthermore, we explored combinational anticancer therapy using HPRP-A1 with the chemical drugs doxorubicin (DOX). The data showed that the anticancer

activity of these drugs against HeLa cell lines was synergistically increased both *in vitro* and *in vivo* [6]. We think this strategy of combination therapy appears to have great clinical potential.

Recently, we are working on the targeting modification of HPRP-A1 and the polymer modification to improve the selectivity and stability. We believe that ACPs as novel anticancer drugs will play an important role for the clinical practices.

References

1. Huang Y (2015) Alpha-helical cationic anticancer peptides: A promising candidate for novel anticancer drugs. *Mini Rev Med Chem* 15: 73-81.
2. Zhao LJ (2013) Comparison on effect of hydrophobicity on the antibacterial and antifungal activities of alpha-helical antimicrobial peptides. *Science China-Chemistry* 56: 1307-1314.
3. Putsep K (1999) Antibacterial peptide from *H. pylori*. *Nature* 398: 671-672.
4. Zhao J (2015) *In vitro* Characterization of the rapid cytotoxicity of anticancer peptide HPRP-A2 through membrane destruction and intracellular mechanism against gastric cancer cell lines. *PLoS One* 10: e0139578.
5. Hao X (2015) TAT Modification of Alpha-helical anticancer peptides to improve specificity and efficacy. *PLoS One* 10: e0138911.
6. Zhao J (2015) Two hits are better than one: Synergistic anticancer activity of α -helical peptides and doxorubicin/epirubicin. *Oncotarget* 6: 1769-1778.