

Orotic Acid: Why it is Important to Understand Its Role in Metabolism?

Thomas P West*, Chunduru J and Eswara C Murahari

Department of Chemistry, Texas A&M University-Commerce, Commerce, USA

The molecule orotic acid was originally thought to be a vitamin that was essential to animal nutrition. Vitamin B₁₃ was originally postulated to be present in distillers dried soluble from grains. It was later determined that vitamin B₁₃ or orotic acid was synthesized by mammals during the synthesis of pyrimidines using the *de novo* pyrimidine biosynthetic pathway [1,2]. In humans and other organisms, orotic acid is synthesized by the enzyme dihydroorotate dehydrogenase which converts dihydroorotate to orotic acid [3]. It has been found that orotic acid improves the metabolism of folic acid and vitamin B₁₂. Orotic acid is found in milk produced by cows and other commercial dairy products derived from milk. The presence of orotate in mammals is important to the development of the central nervous system [1]. The importance of orotic acid in human metabolism can be witnessed in individuals afflicted with orotic aciduria. Orotic aciduria is detected in humans by virtue of affected individuals excreting excess orotic acid. This autosomal recessive disorder has been associated with developmental retardation.

Two forms of orotic aciduria have been distinguished in humans. Type I form of orotic aciduria is caused by the loss of uridine 5'-monophosphate synthase activity. The other form of orotic aciduria is associated with megaloblastic anemia [4-6]. Megaloblastic anemia results due to DNA synthesis inhibition in human red blood cells. Recently, it was noted that the administration of orotic acid in humans improved the condition of hearts with hypertrophy. It was concluded that the positive effect of administering orotic acid to humans with hypertrophic hearts was caused by the release of pyrimidine nucleosides into the blood stream blocking adenine nucleotide depletion in the myocardium [7]. It has also been observed that salts of orotic acid, including zinc or magnesium orotate, can be used in metal ion substitution therapy [5]. Negative effects have also been observed with respect to orotic acid since insulin resistance and hypertension or fatty livers can be induced in rats when orotic acid is included in their diet [8,9]. The hypertension observed in orotic acid-fed rats was thought to be the result of impaired endothelial nitric acid synthesis [8]. With respect to gene transcription, it has been speculated that orotic acid may influence the synthesis of genes involved in cell proliferation in eukaryotes [1]. In prokaryotes, the supplementation of orotic acid to *Pseudomonas fluorescens*, *Pseudomonas putida* and *Pseudomonas lundensis* cells was shown to increase the transcription of the pyrimidine biosynthetic pathway enzyme dihydroorotase, orotate phosphoribosyltransferase or orotidine 5'-monophosphate decarboxylase [10-12].

It is clear that orotic acid can affect gene transcription in both prokaryotes and eukaryotes. In summary, the importance of orotic acid, previously identified as vitamin B13, in metabolism has been known since its discovery in 1905 [1] but recent research findings have demonstrated that orotic acid likely has additional roles in metabolism. Only recently has a more intensive effort been made to learn the relationship between orotic acid levels and its cumulative effect upon metabolism. Further analysis of the role that this pyrimidine base occupies in metabolism needs to be elucidated to more fully understand its overall importance.

References

1. Löffler M, Carrey EA, Zameitat E (2015) Orotic acid, more than just an intermediate of pyrimidine *de novo* synthesis. *J Genet Genomics* 42: 207-219.
2. Löffler M, Carrey EA, Zameitat E (2016) Orotate (orotic acid): An essential and versatile molecule. *Nucleosides Nucleotides Nucleic Acids* 35: 566-577.
3. Garavito MF, Narváez-Ortiz HY, Zimmerman BH (2015) Pyrimidine metabolism: Dynamic and versatile pathways in pathogens and cellular development. *J Genet Genomics* 42: 195-205.
4. Nyhan WL (2005) Disorders of purine and pyrimidine metabolism. *Mol Genet Metab* 86: 25-33.
5. Löffler M, Fairbanks LD, Zameitat E, Marinaki AM, Simmonds HA (2005) Pyrimidine pathways in health and disease. *Trends Mol Med* 11: 430-437.
6. Bailey CJ (2009) Orotic aciduria and uridine monophosphate synthase: A reappraisal. *J Inher Metab Dis* 32: S227-S233.
7. Richards SM, Conyers RAJ, Fisher JL, Rosenfeldt FL (1997) Cardioprotection by orotic acid: Metabolism and mechanism of action. *J Mol Cell Cardiol* 29: 3239-3250.
8. Choi YJ, Yoon Y, Lee KY, Kang YP, Lim DK, et al. (2015) Orotic acid induces hypertension associated with impaired endothelial nitric oxide synthesis. *Toxicol Sci* 144: 307-317.
9. Wang YM, Hu XQ, Xue Y, Li ZJ, Yanagita T, et al (2011) Study on possible mechanism of orotic acid-induced fatty liver in rats. *Nutrition* 27: 571-575.
10. Chu CP, West TP (1990) Pyrimidine biosynthetic pathway of *Pseudomonas fluorescens*. *J Gen Microbiol* 136: 875-880.
11. Santiago MF, West TP (2002) Control of pyrimidine formation in *Pseudomonas putida* ATCC 17536. *Can J Microbiol* 48: 1076-1081.
12. West TP (2009) Regulation of pyrimidine formation in *Pseudomonas lundensis*. *Can J Microbiol* 55: 261-268.

*Corresponding author: Thomas P West, Department of Chemistry, Texas A&M University-Commerce, Commerce, TX 75429-3011, USA, Tel: +1-903-886-5381; Fax: 903-468-6020; E-mail: Thomas.West@tamuc.edu

Received April 22, 2017; Accepted April 24, 2017; Published April 29, 2017

Citation: West TP, Chunduru J, Murahari EC (2017) Orotic Acid: Why it is Important to Understand Its Role in Metabolism? *Biochem Physiol* 6: e157. doi: [10.4172/2168-9652.1000e157](https://doi.org/10.4172/2168-9652.1000e157)

Copyright: © 2017 West TP, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.