

## Ovarian Tissue Transplantation versus Follicular Culture

Laleh Hosseini<sup>1,2</sup>, Mohammad Mehdi Akhond<sup>1</sup>, Mohammad Mehdi Naderi<sup>1</sup>, Banafsheh Heidari<sup>1</sup> and Abolfazl Shirazi<sup>1\*</sup>

<sup>1</sup>Reproductive Biotechnology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran

<sup>2</sup>Research Institute of Animal Embryo Technology, Shahrekord University, Shahrekord, Iran

### Abstract

The banking of ovarian tissue containing large number of primordial follicles has become widespread as one of the promising fertility preservation options in young cancerous patients. There are substantially three approaches in female fertility restoration including embryo and oocyte freezing as well as ovarian tissue cryopreservation. Two former methods require ovarian hyperstimulation which in turn have their own side effects and limitations. The third option as the only feasible approach in prepubertal girls allows immediate cancer treatment. In third method, two main approaches are proposed. One is ovarian tissue transplantation which may not be an applicable approach for all patients especially where there is a chance of reintroducing malignant cells. The second approach with greater interest is one or two steps ovarian follicular culture followed by in vitro maturation of resulting oocytes and the subsequent IVF and IVC procedures. Among various studies conducted in different species, following follicle culture, the live birth has been achieved only in mouse. In this review various aspects of the studies done in this area would be challenged.

**Keywords:** Follicle culture; Ovarian tissue culture; Primordial follicle; Autotransplantation; Three dimensional culture

### Introduction

Fertility preservation provides the pregnancy chances at the right time in patients or those who want to postpone childbearing for social or financial reasons. The majority of patients who can benefit from fertility preservation techniques are cancerous patients. Irreversible follicular and oocyte damage due to chemotherapy and radiotherapy led to the fertility loss in women so that 50% of primordial follicles could be destroyed following radiation [1]. There are several approaches for fertility preservation in female cancerous patients including oocyte or embryo cryopreservation and ovarian tissue banking followed by tissue transplantation or one or two steps follicular culture [2-4].

Embryo cryopreservation has been widely used in many species [5-8] and the oocytes cryopreservation as an alternative method is accounted for more than 200 live births [9-12]. Application of oocyte cryopreservation, however, needs to be more improved in animal species [13]. In prepubertal girls or cancerous patients where the future fertility is desired, the ovarian tissue cryopreservation is more promising. The human ovarian cortex contains the vast majority of the follicular reserve that is less susceptible to cryodamage. Cortical ovarian tissue can orthotopically or heterotopically be autotransplanted and also can be used for follicle culture. In in vitro culture of follicles both follicle development and oocyte health supported during long-term culture [4].

Considering the importance of female fertility preservation in patients at risk of compromised fertility, this mini-review will discuss two main approaches proposed in cancerous patients.

### Transplantation of Ovarian Tissue

#### Autotransplantation

In cancerous patient, when there is no risk of ovarian metastatic involvement, ovarian tissue can be transplanted otherwise an alternative approach should be considered such as one or two step follicular culture. There are mainly two strategies including transplantation of whole ovary or cortical ovarian tissue reimplantation.

In whole ovary transplantation, despite reduction in ischemia and prolonged graft longevity, there are difficulties in cryopreservation and

supplying the required nutrients to deeper parts considering the size of ovary. Nonetheless, transplantation of frozen-thawed whole ovaries in sheep and subsequent oocyte aspiration has resulted in embryo development up to the 8-cell stage [14]. Transplantation of whole cryopreserved ovaries with microanastomosis of the ovarian vascular pedicle resulted in pregnancy and live birth in mouse and sheep [15,16]. The whole human ovary transplantation between monozygotic twins who were discordant for polycystic ovarian failure led to the birth of healthy baby [17].

In second strategy, the cortical ovarian tissue has been successfully transplanted into mouse, sheep, and monkey [18-20]. The first successful human ovarian tissue transplantation was performed between monozygotic twins in 2005 [21]. The results of ovarian cortex reimplantation were more promising as such ovarian activity restoration was observed 3.5 months after reimplantation and successful pregnancy was achieved 9 month after orthotopic reimplantation of ovarian cortex [22]. Ovarian cortex autotransplantation to a peritoneal pocket in the broad ligament led to ovarian function recovery 24 weeks after transplantation and normal pregnancy following the fifth stimulation attempt [23]. So far, 30 live births have been reported after orthotopic reimplantation of cryopreserved ovarian tissues [24]. On the other side the heterotopic transplantation has resulted in production of a four-cell embryo leading to one pregnancy [25].

#### Xenotransplantation

Since in human autotransplantation there are some limitations in assessment of physiological aspects of follicular development, xenotransplantation of ovarian tissue to immunodeficient animals

\*Corresponding author: Abolfazl Shirazi, Reproductive Biotechnology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran, Tel: 00989131821238; E-mail: [A.Shirazi@avicenna.ac.ir](mailto:A.Shirazi@avicenna.ac.ir)

Received June 23, 2014; Accepted July 29, 2014; Published July 31, 2014

Citation: Hosseini L, Akhond MM, Naderi MM, Heidari B, Shirazi A (2014) Ovarian Tissue Transplantation versus Follicular Culture. JFIV Reprod Med Genet 2: 129. doi:10.4172/jfiv.1000129

Copyright: © 2014 Hosseini L, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

provides the opportunity to perform such studies on survival, morphology and functional recovery of ovarian follicles.

The oocytes in bovine secondary follicles grown in Severe Combined Immune Deficiency mice (SCID) were able to resume meiosis and progress to the second metaphase [26]. Xenografting of pig vitrified-warmed and fresh ovarian tissues in SCID mice led to primordial follicles development to the secondary and antral stages, respectively, after two months. In between, though the primordial follicles could maintain their developmental competence after vitrification and warming, their developmental rate was slower than that of fresh counterparts [27]. Fresh and frozen-thawed human ovarian tissue xenograft to SCID mice has increased the proportion of growing follicles as well as the growth of follicle to the antral stage. There was, however, no significant difference in oocyte or follicle diameters between fresh and frozen-thawed tissue grafts. Theca layer in antral follicles of frozen-thawed grafted tissue, however, was significantly thinner than fresh tissue [28].

Xenografting of vitrified-warmed bovine follicles into SCID mice has led to the growth of primordial and secondary follicles to the antral stage, indicating the developmental ability of follicles after vitrification [29]. The survival and development of xenografted human follicles into antral stages after six months was indicated the capacity of oocytes to survive for long term [30]. Xenotransplantation of fresh isolated human follicles into nude mice resulted in primordial follicles activation and formation of follicles surrounded by stroma-like tissue of human origin [31]. Bovine isolated preantral follicles embedded in granulosa and stroma cells matrix were able to survive and grow 14 days after renal subcapsular xenotransplantation. This matrix provides an *in vivo* model to study preantral follicles development [32]. Observation of infiltrating blood capillaries after xenograft of an alginate-matrigel matrix containing isolated ovarian cells was another promising event in developing a biodegradable scaffold [33].

One major limitation in grafted ovarian tissue is a considerable follicles loss. Ischemia and the consequent apoptosis that occurs before efficient revascularization is the cause of this reduction, as >50% of primordial follicles are lost following ovarian transplantation. The loss of follicles may also be due to premature activation of the transplanted follicular pool [31]. Lack of antimullerian hormone in ovarian grafts is proposed for premature follicular activation [34]. Graft pretreatment with vascular endothelial growth factor A (VEGF-A) and vitamin E besides host treatment with vitamin E and gonadotropins could improve the survival of grafted human ovarian tissue by reducing apoptosis [35]. Additionally, the ischemic injury was decreased by sphingosine-1-phosphate supplementation through acceleration of angiogenic process and reduction of tissue hypoxia [36]. VEGF and fibroblast growth factor b (bFGF), especially in combination, through triggering angiogenesis and reducing apoptosis could increase the survival of transplanted human ovarian tissue [37].

## Follicle Culture

### Ovarian tissue culture

Since the early 1990s, ovarian follicle culture techniques have been developed with the aim of achieving competent oocytes. In some species such as mouse the culture of ovarian follicles in the form of whole ovary culture is executable. Moreover, the small size and soft texture of the ovaries provide the opportunity to achieve intact follicles following enzymatic digestion. In contrast, in human and most domestic species, the ovaries are too large to be organ-cultured, and the toughness of ovarian stroma impairs (compromises) the follicle intactness following enzymatic dissociation. To conquer these problems, an *in vitro* system

was developed for culturing small pieces of ovarian cortex prior to follicle isolation. Cortical pieces were cultured in medium supplemented with serum plus insulin, transferrin, and selenium (ITS) where the primordial follicles were activated to become primary follicles. Though, in ovarian cortical strip culture, the follicles were deprived from the *in vivo* endocrine and paracrine factors, the follicles were received the effects of follicle-stromal interactions. Removing stromal cells and culturing the flattened tissues led to the greater activation and faster follicles growth [38].

### Hormones and growth factors in tissue culture

Studies on animal models considering the effects of hormones and growth factors on follicular development and survival have widely provided a basis for human follicle culture experiments.

In human ovarian tissue culture, ascorbic acid and cyclic adenosine monophosphate have been used to prevent apoptosis [39]. The positive effects of insulin growth factor I and II (IGF-I and II) in reducing atresia, increasing growing follicles and follicular integrity has been established [40]. There are also evidences indicating the promotory effect of FGF on human follicle development as well as the stimulatory effect of growth and differentiation factor 9 (GDF9) on follicle integrity and primordial follicle activation [41,42]. In between, anti mullerian hormone has an inhibitory effect on human ovarian follicular development by suppressing primordial follicle activation [43].

Studies in animal have indicated the positive effect of GDF9 on ultrastructural integrity of goat preantral follicles and primordial follicles activation [44]. The presence of follicle stimulating hormone (FSH) and FGF-2 in ovarian tissue culture has shown the promotory effects of FSH on percentage of the primary follicles. FSH alone or combined with FGF-2 increased the growth and integrity of caprine preantral follicle [45]. In cattle, FSH in combination of GDF-9 or bFGF has increased the rate of normal follicles and decreased the rate of apoptotic cells [46]. Presence of indol acetic acid (IAA), EGF, and FSH has maintained ultrastructural integrity of sheep primordial follicles and oocytes [47]. In mice, culture of prepubertal ovarian explants with R-spondin2, stem cell growth factor, has promoted primary follicles activation [48].

### Isolated follicle culture

In culturing ovarian tissues, as the follicles develop to the secondary stage the cortical tissue environment becomes inhibitory to further growth. Therefore, ovarian tissue culture cannot support follicle development to the more advanced stages so that a multi-step culture system is required to support further development. Ovarian follicles can be mechanically or enzymatically isolated from the cortex and then cultured for further development [38]. In human, considering the toughness of ovarian cortex, mechanical isolation of intact primordial follicles is difficult and enzymatic digestion is more appropriate.

For isolated follicles culture, two approaches have been proposed including two- and three-dimensional culture systems. In two-dimensional culture system, referred to attached follicle approach, concurrent with follicular development the proliferating granulosa cells attach to the plate and migrate away from the oocyte. Therefore, the granulosa cells are not able to support properly the follicle development because of their spatial disarrangement. In contrast, in three dimensional culture system, the follicle is able to maintain its three-dimensional architecture as the follicular growth occurs radially from the center of the follicle. This structure provides mechanical support which is essential for maintaining cell-cell contacts and paracrine signaling [49].

There are various materials to maintain 3D architecture such as natural (e.g. collagen) and synthetic (e.g. alginate) hydrogels. Alginate encapsulation due to its flexibility and partial rigidity can mimic the extracellular matrix. This 3D architecture besides facilitating molecular exchange between the follicle and the culture medium can provide the proper conditions for cell proliferation and antrum formation [50,51].

### Two dimensional follicle culture

Although this system has been successful in production of live murine pups, it has not been successful in bovine, ovine, and human species. In human and large animals to maintain the communication between the oocyte and granulosa cells, which is necessary for follicular development, is more difficult due to considerable size of follicle and the longer time required to culture. Two-dimensional culture of human follicles has shown the high level of follicular atresia due to loss of connections between the oocyte and the granulosa cells [52,53]. In mouse, two-step culture system, including an 8-day culture of whole newborn ovaries followed by culture of isolated secondary follicles and the subsequent culture of obtained oocytes has resulted in production of normal offspring [54].

### Three dimensional follicle culture

**Experience with primordial and primary follicles:** The two-step primordial and primary follicles culture in serum free media covering 6 days culture of human ovarian cortical strip followed by 4 days culture of isolated follicles resulted in formation of antrum in cultured follicles [38].

In 3-dimensional culture system, among different contributing factors, the type and rigidity of follicular extracellular matrix by maintaining the connection between oocyte and granulosa cells and regulation of numerous cellular processes has an important role in follicle development. Seven days culture of frozen-thawed human primordial follicles in alginate hydrogel 1% led to an increase in follicle size with survival rate of 90% [55]. Long term culture of ovine primary follicles in fibronectin-coated wells in serum-free medium resulted in follicular progression to the secondary follicle stage [56]. Encapsulating the macaque primary and secondary follicles in alginate and fibrin alginate promoted the follicle development into antral stage. In between, the culture of secondary follicle in alginate yielded an MII oocyte which after fertilization could further develop to morula stage. Depends on developmental stage of follicle, the type of scaffold has an important effect on subsequent follicle development [57]. Culturing macaque primordial follicles in different concentrations of alginate demonstrated 0.5% alginate could maintain the integrity of cultured follicles for up to 3 days, while the integrity was lost after 6 days. In contrast, follicular integrity could be maintained in 2% alginate up to sixth day. Therefore, in order to culture macaque primordial follicles, a more rigid environment was needed to support follicle growth in vitro [58].

In between, hormones and growth factors through endocrine, paracrine, and autocrine mechanisms have major role in follicle development. In bovine, primary follicles culture in presence of gonadotropins and growth factors during 21 days has optimized the culture system to support follicle growth to antral stage [59]. Considering to the paracrine mechanism, it would be expected that group culture of follicles could better support the follicle growth compared to single culture system. In vitro culture of sheep primordial follicles in higher densities (50 or 100 lectin-aggregated follicles per well) has significantly increased the diameter and survival of oocytes and induced granulosa cell differentiation [60]. In mouse, group culture of primary follicles has

increased the growth and survival of follicles leading to antral follicles formation containing meiotically competent oocytes [61].

Any follicular growth system should be able to promote quiescent primordial follicles to grow within ovarian cortical tissue. To this end, and considering the negative regulatory effects of PTEN and FOXO3, as inhibitors of primordial follicle activation, application of their inhibitors may promote the growth of human and mouse follicular reserves [62,63]. It has also been established that inhibition of mTOR, mammalian target of rapamycin, may decrease human primordial follicles activation, leading to oocyte loss in growing follicles [64].

**Experience with preantral follicle:** Until recently the work has focused on secondary follicles which are less abundant in adult ovaries. Factors controlling in vitro follicular development are including extra cellular matrix (ECM) and culture condition that regulate numerous cellular process.

Structural and biochemical design of the ECM has an influence on growth of follicles. Alginate has been used successfully in preantral follicle culture. Culture of mouse encapsulated secondary follicles within alginate matrix could support follicle development and resulted in oocyte maturation required for fertilization and live birth [65,66]. Antral follicles have also been achieved through 30 days culture of human fresh secondary follicles in alginate in the presence of FSH [50]. Macaque secondary follicle culture in alginate yielded an MII oocyte that following IVF could cleave and reach the morula stage [57].

Fibrin alginate can also be used in preantral follicle culture. As the follicle expands it can produce proteases which in turn through fibrin degradation lead to the reduction of compressive force and support volume increase concurrent with follicular growth. Culture of mouse secondary follicles in alginate and fibrin alginate showed that the rate of meiotically competent oocytes produced by culture in fibrin alginate was greater than alginate alone [51]. The progression of mouse follicles to antral stage in fibrin alginate after aprotinin removal indicate that delay in fibrin degradation by protease inhibitors may be suitable for smaller follicles that require longer culture time [67]. Semidegradable fibrin-alginate matrices allowed the growth of baboon preantral follicle to the antral stage in an FSH-independent manner [68]. Apart from commonly used matrix (alginate), the application of hyaluronan, in mouse, could increase the survival rate and germinal vesicle breakdown [69].

In follicle culture, the ovarian cycle stages during which follicles are collected as well as the physical properties of the matrix are important. Culture of monkey secondary follicles throughout 30 days showed the higher survival rate and growth rate in follicles obtained from prepubertal and adult monkeys, respectively [70]. In another study in monkey, culture of follicles in alginate scaffold, obtained during early follicular phase had a higher survival rate than those collected during the luteal phase [71]. In mouse, 0.25% and 0.5% alginate could better support the growth of follicles and antrum formation compared to the higher alginate concentrations despite the lack of difference in survival rates and oocyte developmental competence [72].

**Hormones and growth factors in preantral follicle culture:** Besides the impact of used system on follicular culture, the contents of culture media including gonadotropins and growth factors are effective in follicular development.

Concerning the effects of gonadotropins on follicular culture, 6-days culture of sheep follicles in the presence of FSH and thyroxin has greatly improved the proportion of matured oocyte [73]. In prepubertal monkey, the culture of follicles in presence of LH after 30

days had a positive effect on follicle diameter [70]. Interestingly, while the application of FSH, alone, had a positive effect on follicular growth, medium supplementation with both FSH and LH had a lower effect [71]. No follicles were survived after forty days culture of alginate-encapsulated primate secondary follicles in the absence of rhFSH [74]. In macaque, the highest survival rate of secondary follicles was observed in presence of high or medium FSH concentrations. FSH had different effect on follicular steroidogenesis based on its concentration. While, steroid production by growing follicles was stimulated in the presence of high FSH concentrations, this production at low FSH concentrations was promoted by LH [75]. In baboon, FSH even had a negative influence on preantral follicular health by disrupting the integrity of oocyte and cumulus cells connections so that the growing follicles could produce MII oocytes with normal spindle structure in the absence of FSH [68]. From above, it could be concluded that despite the positive effects of FSH on preantral follicular development, in many species, there is an exception so that in baboon, it had a negative effect on preantral follicular growth.

In caprine, while the growth of both small and large follicles was stimulated by the presence of growth factors, FSH could stimulate only the development of small but not large preantral follicles [76]. The culture of buffalo preantral follicles in the presence of growth factors and FSH resulted in follicle survival for up to 20 days and early antrum formation. In meantime, IGF-I supplementation had a significantly positive effect on growth and survival of cultured follicles. However, the results were inverted when IGF-I was accompanied with bFGF [77]. In sheep, supplementation of preantral follicles culture medium with growth factors and hormones (ITS, IGF-I, insulin and GH) could properly support follicular development so that IGF-I in combination with GH had the best effect. Though, the cultured oocytes in any concentration of TGF failed to develop to MII stage [78]. Human isolated secondary follicles undergo differentiation after a 4 days in vitro culture in the presence of activin [38]. In feline, 14-days culture of follicles in the presence of activin A promoted granulosa cell proliferation and preantral follicles growth and viability whereas no beneficial effects was observed by thyroxin supplementation [79]. There is also evidence indicating the effect of nitric oxide on preantral follicle culture. As shown, nitric oxide depends on its concentration could play a dual role on follicle growth and survival, whereas the lower doses could stimulate the follicle survival, growth, and antrum formation, the higher concentrations had an inhibitory effects [80]. Moreover, group culture of goat preantral follicles, increased the rates of follicular survival and growth as well as the number of grown oocytes and meiosis resumption [81].

Besides the content of culture medium, the culture condition, e.g. oxygen tension, has an effect on follicle development. It was found that 20% O<sub>2</sub> was more efficient than 5% in goat preantral follicular survival and growth and resumption of oocyte meiosis throughout 30 days culture [82]. In macaque, the effects of various concentrations of fetuin and O<sub>2</sub> on encapsulated secondary follicle showed the highest follicular survival rate in the presence of 1 mg/ml fetuin at 5% O<sub>2</sub> [75].

### Follicle co-culture

Co-culturing of buffalo preantral follicles with somatic cells led to the higher growth rate and survivability [83]. In mouse, follicles co cultured with stromal cells grew more with the greater survival rate [84]. Co-culture of mouse alginate encapsulated follicles with mouse embryonic fibroblast (MEF) promoted the growth of secondary and primary follicles to antral stage after 14 days while the follicles were degenerated, within 6-10 days, in the absence of fibroblasts. In between,

survival rate of 100-µm follicles was significantly higher than 70-µm follicles [85]. The presence of MEF and media supplementation with activin A especially when used together had a positive effect on growth, survivability, and hormonal production of preantral follicles. However, no significant differences were observed in antrum formation, ovulation rate, and subsequent embryonic development [86].

**Follicle development evaluations:** There are numerous criteria which might be considered in assessment of follicular development including follicle survival and growth, steroidogenesis, production of paracrine/autocrine factors, the ability of oocyte to mature, and the pattern of genes expression. Follicle survivability can be identified through assessment of basement membrane intactness, presence of granulosa cells, and the presence of round and centrally located oocyte with a visible zona pellucida. The in vitro follicular production of endocrine factors such as progesterone (P4), androstenedione (A4), and estradiol (E2), as well as paracrine/autocrine factors, such as anti-Müllerian hormone (AMH) and vascular endothelial growth factor (VEGF) that positively correlates with follicle growth and development, might be considered as other criteria. The nuclear maturational status of oocytes derived from in vitro antral follicles following 34 h hCG treatment can be considered as another criterion. In performed studies, no difference was observed in gene expression between the in vitro and in vivo-developed antral follicles in major steroidogenic enzymes except for the upregulation of low density lipoprotein receptor gene that might be related to the prolonged exposure to exogenous FSH. The VEGF downregulation and anti-apoptotic factors upregulation in in vitro-developed primate antral follicles compared to in vivo-derived ones might indicate the influence of culture conditions on gene expression [87].

### Conclusion

The application of ovarian tissue grafting or culture of isolated follicles followed by in vitro production of embryos has their own benefits and defects. Nonetheless, autotransplantation of thawed ovarian tissue has been the only method that resulted in birth of healthy baby. The major drawback of this method, probable re-introduction of cancer cells, has forced the investigators to think of other methods such as multi-steps ovarian follicular culture. In this method despite the remarkable reduction in risk of cancer cells transmission, because of our inadequate knowledge about follicular development and its complexity, no proper system has been fully optimized to meet all challenges posed to extended human follicle growth in vitro. Therefore an introduction of a practical and reliable method has remained to be further investigated.

### References

1. Wallace WH, Thomson AB, Kelsey TW (2003) The radiosensitivity of the human oocyte. *Hum Reprod* 18: 117-121.
2. Salama M, Winkler K, Murach KF, Seeber B, Ziehr SC, et al. (2013) Female fertility loss and preservation: threats and opportunities. *Ann Oncol* 24: 598-608.
3. Telfer EE, Zelinski MB (2013) Ovarian follicle culture: advances and challenges for human and nonhuman primates. *Fertil Steril* 99: 1523-1533.
4. Smits J, Dolmans MM, Donnez J, Fortune JE, Hovatta O, et al. (2010) Current achievements and future research directions in ovarian tissue culture, in vitro follicle development and transplantation: implications for fertility preservation. *Hum Reprod Update* 16: 395-414.
5. Roque M, Lattes K, Serra S, Solà I, Geber S, et al. (2013) Fresh embryo transfer versus frozen embryo transfer in in vitro fertilization cycles: a systematic review and meta-analysis. *Fertil Steril* 99: 156-162.
6. Shirazi A, Soleimani M, Karimi M, Nazari H, Ahmadi E, et al. (2010) Vitrification of in vitro produced ovine embryos at various developmental stages using two methods. *Cryobiology* 60: 204-210.

7. Shirazi A, Nazari H, Ahmadi E, Heidari B, Shams-Esfandabadi N (2009) Effect of culture system on survival rate of vitrified bovine embryos produced in vitro. *Cryobiology* 59: 285-290.
8. Heo YT, Lee SE, Jang WI, Min SG, et al. (2013) A new modified cut standard straw vitrification technique reduces the apoptosis of mouse blastocysts and generates more live mouse offspring. *Cryo Letters* 34: 598-607.
9. Garcia-Velasco JA, Domingo J, Cobo A, Martínez M, Carmona L, et al. (2013) Five years' experience using oocyte vitrification to preserve fertility for medical and nonmedical indications. *Fertil Steril* 99: 1994-1999.
10. Kim MK, Lee DR, Han JE, Kim YS, Lee WS, et al. (2011) Live birth with vitrified-warmed oocytes of a chronic myeloid leukemia patient nine years after allogeneic bone marrow transplantation. *J Assist Reprod Genet* 28: 1167-1170.
11. Vajta G, Rienzi L, Cobo A, Yovich J (2010) Embryo culture: can we perform better than nature? *Reprod Biomed Online* 20: 453-469.
12. Gosden RG (2005) Prospects for oocyte banking and in vitro maturation. *J Natl Cancer Inst Monogr* : 60-63.
13. Shirazi A, Taheri F, Nazari H, Norbakhsh-Nia M, Ahmadi E, et al. (2012) Developmental competence of ovine oocyte following vitrification: effect of oocyte developmental stage, cumulus cells, cytoskeleton stabiliser, FBS concentration, and equilibration time. *Zygote* 14: 1-9.
14. Arav A, Revel A, Nathan Y, Bor A, Gacitua H, et al. (2005) Oocyte recovery, embryo development and ovarian function after cryopreservation and transplantation of whole sheep ovary. *Hum Reprod* 20: 3554-3559.
15. Wang X, Chen H, Yin H, Kim SS, Lin Tan S, et al. (2002) Fertility after intact ovary transplantation. *Nature* 415: 385.
16. Imhof M, Bergmeister H, Lipovac M, Rudas M, Hofstetter G, et al. (2006) Orthotopic microvascular reanastomosis of whole cryopreserved ovine ovaries resulting in pregnancy and live birth. *Fertil Steril* 85 Suppl 1: 1208-1215.
17. Silber SJ, Grudzinskas G, Gosden RG (2008) Successful pregnancy after microsurgical transplantation of an intact ovary. *N Engl J Med* 359: 2617-2618.
18. Candy CJ, Wood MJ, Whittingham DG (2000) Restoration of a normal reproductive lifespan after grafting of cryopreserved mouse ovaries. *Hum Reprod* 15: 1300-1304.
19. Bordes A, Lornage J, Demirci B, Franck M, Courbiere B, et al. (2005) Normal gestations and live births after orthotopic autograft of vitrified-warmed hemiovaries into ewes. *Hum Reprod* 20: 2745-2748.
20. Lee DM, Yeoman RR, Battaglia DE, Stouffer RL, Zelinski-Wooten MB, et al. (2004) Live birth after ovarian tissue transplant. *Nature* 428: 137-138.
21. Silber SJ, Lenahan KM, Levine DJ, Pineda JA, Gorman KS, et al. (2005) Ovarian transplantation between monozygotic twins discordant for premature ovarian failure. *N Engl J Med* 353: 58-63.
22. Donnez J, Squiffet J, Jadoul P, Demylle D, Cheron AC, et al. (2011) Pregnancy and live birth after autotransplantation of frozen-thawed ovarian tissue in a patient with metastatic disease undergoing chemotherapy and hematopoietic stem cell transplantation. *Fertil Steril* 95: 1787e1-1787e4.
23. Donnez J, Jadoul P, Pirard C, Hutchings G, Demylle D, et al. (2012) Live birth after transplantation of frozen-thawed ovarian tissue after bilateral oophorectomy for benign disease. *Fertil Steril* 98: 720-725.
24. Donnez J, Dolmans MM, Pellicer A, Diaz-Garcia C, Sanchez Serrano M, et al. (2013) Restoration of ovarian activity and pregnancy after transplantation of cryopreserved ovarian tissue: a review of 60 cases of reimplantation. *Fertil Steril* 99: 1503-1513.
25. Stern CJ, Gook D, Hale LG, Agresta F, Oldham J, et al. (2013) First reported clinical pregnancy following heterotopic grafting of cryopreserved ovarian tissue in a woman after a bilateral oophorectomy. *Human Reproduction* 28: 2996-2999.
26. Senbon S, Ota A, Tachibana M, Miyano T (2003) Bovine oocytes in secondary follicles grow and acquire meiotic competence in severe combined immunodeficient mice. *Zygote* 11: 139-149.
27. Moniruzzaman M, Bao RM, Taketsuru H, Miyano T (2009) Development of vitrified porcine primordial follicles in xenografts. *Theriogenology* 72: 280-288.
28. Amorim CA, David A, Dolmans MM, Camboni A, Donnez J, et al. (2011) Impact of freezing and thawing of human ovarian tissue on follicular growth after long-term xenotransplantation. *J Assist Reprod Genet* 28: 1157-1165.
29. Bao RM, Yamasaka E, Moniruzzaman M, Hamawaki A, Yoshikawa M, et al. (2010) Development of vitrified bovine secondary and primordial follicles in xenografts. *Theriogenology* 74: 817-827.
30. Dolmans MM, Yuan WY, Camboni A, Torre A, Van Langendonck A, et al. (2008) Development of antral follicles after xenografting of isolated small human preantral follicles. *Reprod Biomed Online* 16: 705-711.
31. Dolmans MM, Martínez-Madrid B, Gadisseux E, Guiot Y, Yuan WY, et al. (2007) Short-term transplantation of isolated human ovarian follicles and cortical tissue into nude mice. *Reproduction* 134: 253-262.
32. Aerts JM, Martínez-Madrid B, Leroy JL, Van Aelst S, Bols PE (2010) Xenotransplantation by injection of a suspension of isolated preantral ovarian follicles and stroma cells under the kidney capsule of nude mice. *Fertil Steril* 94: 708-714.
33. Vanacker J, Luyckx V, Dolmans MM, Des Rieux A, Jaeger J, et al. (2012) Transplantation of an alginate-matrigel matrix containing isolated ovarian cells: first step in developing a biodegradable scaffold to transplant isolated preantral follicles and ovarian cells. *Biomaterials* 33: 6079-6085.
34. David A, Van Langendonck A, Gilliaux S, Dolmans MM, Donnez J, et al. (2012) Effect of cryopreservation and transplantation on the expression of kit ligand and anti-Müllerian hormone in human ovarian tissue. *Hum Reprod* 27: 1088-1095.
35. Abir R, Fisch B, Jessel S, Felz C, Ben-Haroush A, et al. (2011) Improving posttransplantation survival of human ovarian tissue by treating the host and graft. *Fertil Steril* 95: 1205-1210.
36. Soleimani R, Heytens E, Oktay K (2011) Enhancement of neoangiogenesis and follicle survival by sphingosine-1-phosphate in human ovarian tissue xenotransplants. *PLoS One* 6: e19475.
37. Wang L, Ying YF, Ouyang YL, Wang JF, Xu J (2013) VEGF and bFGF increase survival of xenografted human ovarian tissue in an experimental rabbit model. *J Assist Reprod Genet* 30: 1301-1311.
38. Telfer EE, McLaughlin M, Ding C, Thong KJ (2008) A two-step serum-free culture system supports development of human oocytes from primordial follicles in the presence of activin. *Hum Reprod* 23: 1151-1158.
39. Scott J, Zhang P, Hreinsson J, Hovatta O (2004) Cyclic guanosine monophosphate improves the survival of human ovarian follicles in culture. *Reproductive Biomedicine Online* 8: 319-324.
40. Fabbri R, Pasquinelli G, Montanaro L, Mozzanega B, Magnani V, et al. (2007) Healthy early preantral follicle can be obtained in a culture of frozen-thawed human ovarian tissue of 32 weeks. *Ultrastruct Pathol* 31: 257-262.
41. Garor R, Abir R, Erman A, Felz C, Nitke S, et al. (2009) Effects of basic fibroblast growth factor on in vitro development of human ovarian primordial follicles. *Fertil Steril* 91: 1967-1975.
42. Kedem A, Fisch B, Garor R, Ben-Zaken A, Gizunterman T, et al. (2011) Growth differentiating factor 9 (GDF9) and bone morphogenetic protein 15 both activate development of human primordial follicles in vitro, with seemingly more beneficial effects of GDF9. *J Clin Endocrinol Metab* 96: 1246-1254.
43. Carlsson IB, Scott JE, Visser JA, Ritvos O, Themmen AP, et al. (2006) Anti-Müllerian hormone inhibits initiation of growth of human primordial ovarian follicles in vitro. *Hum Reprod* 21: 2223-2227.
44. Martins FS, Celestino JJ, Saraiva MV, Matos MH, Bruno JB, et al. (2008) Growth and differentiation factor-9 stimulates activation of goat primordial follicles in vitro and their progression to secondary follicles. *Reprod Fertil Dev* 20: 916-924.
45. Matos MH, Bruno JB, Rocha RM, Lima-Verde IB, Santos KD, et al. (2011) In vitro development of primordial follicles after long-term culture of goat ovarian tissue. *Res Vet Sci* 90: 404-411.
46. Tang K, Yang WC, Li X, Wu CJ, Sang L, et al. (2012) GDF-9 and bFGF enhance the effect of FSH on the survival, activation, and growth of cattle primordial follicles. *Anim Reprod Sci* 131: 129-134.
47. Andrade ER, Maddox-Hyttel P, Landim-Alvarenga Fda C, Viana Silva JR, Alfieri AA, et al. (2010) Ultrastructure of Sheep Primordial Follicles Cultured in the Presence of Indol Acetic Acid, EGF, and FSH. *Vet Med Int* 2011: 670987.
48. Cheng Y, Kawamura K, Takae S, Deguchi M, Yang Q, et al. (2013) Oocyte-derived R-spondin2 promotes ovarian follicle development. *FASEB J* 27: 2175-2184.
49. West ER, Shea LD, Woodruff TK (2007) Engineering the follicle microenvironment. *Semin Reprod Med* 25: 287-299.
50. Xu M, Barrett SL, West-Farrell E, Kondapalli LA, Kiesewetter SE, et al. (2009)

- In vitro grown human ovarian follicles from cancer patients support oocyte growth. *Hum Reprod* 24: 2531-2540.
51. Shikanov A, Xu M, Woodruff TK, Shea LD (2009) Interpenetrating fibrin-alginate matrices for in vitro ovarian follicle development. *Biomaterials* 30: 5476-5485.
52. Roy SK, Treacy BJ (1993) Isolation and long-term culture of human preantral follicles. *Fertil Steril* 59: 783-790.
53. Abir R, Franks S, Mobberley MA, Moore PA, Margara RA, et al. (1997) Mechanical isolation and in vitro growth of preantral and small antral human follicles. *Fertil Steril* 68: 682-688.
54. O'Brien MJ, Pendola JK, Eppig JJ (2003) A revised protocol for in-vitro development of mouse oocytes from primordial follicles dramatically improves their developmental competence. *Biol Reprod* 68: 1682-1686.
55. Amorim CA, Van Langendonck A, David A, Dolmans MM, Donnez J (2009) Survival of human pre-antral follicles after cryopreservation of ovarian tissue, follicular isolation and in-vitro culture in a calcium alginate matrix. *Human Reproduction* 24: 92-99.
56. Muruvi W, Picton HM, Rodway RG, Joyce IM (2009) In vitro growth and differentiation of primary follicles isolated from cryopreserved sheep ovarian tissue. *Anim Reprod Sci* 112: 36-50.
57. Xu J, Lawson MS, Yeoman RR, Molskness TA, Ting AY, et al. (2013) Fibrin promotes development and function of macaque primary follicles during encapsulated three-dimensional culture. *Hum Reprod* 28: 2187-2200.
58. Hornick JE, Duncan FE, Shea LD, Woodruff TK (2012) Isolated primate primordial follicles require a rigid physical environment to survive and grow in vitro. *Hum Reprod* 27: 1801-1810.
59. Sun J, Li X (2013) Growth and antrum formation of bovine primary follicles in long-term culture in vitro. *Reprod Biol* 13: 221-228.
60. Muruvi W, Picton HM, Rodway RG, Joyce IM (2005) In vitro growth of oocytes from primordial follicles isolated from frozen-thawed lamb ovaries. *Theriogenology* 64: 1357-1370.
61. Hornick JE, Duncan FE, Shea LD, Woodruff TK (2013) Multiple follicle culture supports primary follicle growth through paracrine-acting signals. *Reproduction* 145: 19-32.
62. Li J, Kawamura K, Cheng Y, Liu S, Klein C, et al. (2010) Activation of dormant ovarian follicles to generate mature eggs. *Proc Natl Acad Sci U S A* 107: 10280-10284.
63. Adhikari D, Gorre N, Risal S, Zhao Z, Zhang H, et al. (2012) The safe use of a PTEN inhibitor for the activation of dormant mouse primordial follicles and generation of fertilizable eggs. *PLoS One* 7: e39034.
64. McLaughlin M, Patrizio P, Kayisli U, Luk J, Thomson TC, et al. (2011) mTOR kinase inhibition results in oocyte loss characterized by empty follicles in human ovarian cortical strips cultured in vitro. *Fertil Steril* 96: 1154-1159.
65. Xu M, Kreeger PK, Shea LD, Woodruff TK (2006) Tissue-engineered follicles produce live, fertile offspring. *Tissue Eng* 12: 2739-2746.
66. Wang X, Catt S, Pangestu M, Temple-Smith P (2011) Successful in vitro culture of pre-antral follicles derived from vitrified murine ovarian tissue: oocyte maturation, fertilization, and live births. *Reproduction* 141: 183-191.
67. Shikanov A, Xu M, Woodruff TK, Shea LD (2011) A method for ovarian follicle encapsulation and culture in a proteolytically degradable 3 dimensional system. *J Vis Exp* .
68. Xu M, Fazleabas AT, Shikanov A, Jackson E, Barrett SL, et al. (2011) In vitro oocyte maturation and preantral follicle culture from the luteal-phase baboon ovary produce mature oocytes. *Biol Reprod* 84: 689-697.
69. Desai I, Abdelhafez F, Calabro A, Falcone T (2012) Three dimensional culture of fresh and vitrified mouse pre-antral follicles in a hyaluronan-based hydrogel: a preliminary investigation of a novel biomaterial for in vitro follicle maturation. *Reprod Biol Endocrinol* 10: 29.
70. Zelinski MB, Bernuci M, Lawson M, Yeoman RR, Stouffer RL (2008) Culture of preantral follicles from prepubertal and adult rhesus monkeys in a three-dimensional (3D) alginate matrix: FSH required for growth to early antral stage. *Fertil Steril* 90: S275.
71. Xu M, West-Farrell ER, Stouffer RL, Shea LD, Woodruff TK, et al. (2009) Encapsulated three-dimensional culture supports development of nonhuman primate secondary follicles. *Biol Reprod* 81: 587-594.
72. Xu M, West E, Shea LD, Woodruff TK (2006) Identification of a stage-specific permissive in vitro culture environment for follicle growth and oocyte development. *Biol Reprod* 75: 916-923.
73. Arunakumari G, Vagdevi R, Rao BS, Naik BR, Naidu KS, et al. (2007) Effect of hormones and growth factors on in vitro development of sheep preantral follicles. *Small Rumin Res* 70: 93-100.
74. Xu J, Bernuci MP, Lawson MS, Yeoman RR, Fisher TE, et al. (2010) Survival, growth, and maturation of secondary follicles from prepubertal, young and older adult, rhesus monkeys during encapsulated three-dimensional (3D) culture: effects of gonadotropins and insulin. *Reproduction* 140: 685-697.
75. Xu J, Lawson MS, Yeoman RR, Zelinski MB, Stouffer RL (2011) Secondary follicle growth and oocyte maturation during encapsulated three-dimensional culture in rhesus monkeys: effects of gonadotrophins, oxygen and fetuin. *Human Reproduction* 26: 1061-1072.
76. Rajarajan K, Rao BS, Vagdevi R, Tamilmani G, Arunakumari G, et al. (2006) Effect of various growth factors on the in vitro development of goat preantral follicles. *Anim Reprod Sci* 63: 204-212.
77. Sharma GT1, Dubey PK, Meur SK (2009) Survival and developmental competence of buffalo preantral follicles using three-dimensional collagen gel culture system. *Anim Reprod Sci* 114: 115-124.
78. Arunakumari G, Shanmugasundaram N, Rao VH (2010) Development of morulae from the oocytes of cultured sheep preantral follicles. *Theriogenology* 74: 884-894.
79. Wongbandue G, Jewgenow K, Chatdarong K (2013) Effects of thyroxin (T4) and activin A on in vitro growth of preantral follicles in domestic cats. *Theriogenology* 79: 824-832.
80. Dubey PK, Tripathi V, Singh RP, Saikumard G, Natha A, et al. (2012) Expression of nitric oxide synthase isoforms in different stages of buffalo (Bubalus bubalis) ovarian follicles: Effect of nitric oxide on in vitro development of preantral follicle. *Theriogenology* 77: 280-291.
81. Duarte AB, Chaves RN, Araújo VR, Celestino JJ, Silva GM, et al. (2011) Follicular interactions affect the in vitro development of isolated goat preantral follicles. *Zygote* 19: 215-227.
82. Silva CM, Matos MH, Rodrigues GQ, Faustino LR, Pinto LC, et al. (2010) In vitro survival and development of goat preantral follicles in two different oxygen tensions. *Anim Reprod Sci* 117: 83-89.
83. Ramesh SH, Gupta P, Nandi S, Manjunatha B (2010) In vitro development of buffalo preantral follicles in co-culture with cumulus and granulosa cells. *Veterinarski Arhiv* 80: 41-50.
84. Tingen C, Kiesewetter SE, Jozefik J, Thomas C, Tagler D, et al. (2011) A macrophage and theca cell-enriched stromal cell population influences growth and survival of immature murine follicles in vitro. *Reproduction* 141: 809-820.
85. Tagler D, Tu T, Smith RM, Anderson NR, Tingen CM, et al. (2012) Embryonic fibroblasts enable the culture of primary ovarian follicles within alginate hydrogels. *Tissue Eng Part A* 8: 1229-1238.
86. Karimpour Malekshah A, Heidari M, Parivar K, Azami NS (2014) The effects of fibroblast co-culture and activin A on in vitro growth of mouse preantral follicles. *Iran Biomed J* 18: 49-54.
87. Xu J, Xu M, Bernuci MP, Fisher TE, Shea LD, et al. (2013) Primate follicular development and oocyte maturation in vitro. *Adv Exp Med Biol* 761: 43-67.

Citation: Hosseini L, Akhond MM, Naderi MM, Heidari B, Shirazi A (2014) Ovarian Tissue Transplantation versus Follicular Culture. JFIV Reprod Med Genet 2: 129. doi:10.4172/jfiv.1000129