

Parameters of Physiological Responses and Meat Quality in Poultry Subjected to Transport Stress

Jing Qi^{1*}, Yaru Zhang², Zixiong Zhou¹ and Umma Habiba¹

¹College of Veterinary Medicine, Chonbuk National University, Iksan, Republic of Korea

²College of Veterinary Medicine, China Agricultural University, Beijing, China

*Corresponding author: Jing Qi, 1College of Veterinary Medicine, Chonbuk National University, Iksan, Republic of Korea, Tel: 01048891681; E-mail: qijingvet@163.com

Received date: February 2, 2017; Accepted date: March 06, 2017; Published date: March 14, 2017

Copyright: © 2017 Qi J, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Abstract

Transport is unavoidable before slaughter in the poultry industry. The birds may be exposed to various stressors during transport, including birds health condition, fasting, handling, (un)loading, crating density, transport time, vehicle design, trailer microclimate and lairage time. To cope with transport stress, poultry develops physiological responses, which may subsequently affect meat quality and mortality. The paper aims to investigate the effects of transport duration, trailer microclimate and loading density on physiological responses and meat quality of birds. Transport is likely to cause high stress level and energy consumption in poultry according to physiological parameters. The impacts of transport stressors on meat quality are inconsistent or even contradictory. To get consistent results of meat quality following transport, unified measurement methods and the transport conditions are demanded. Furthermore, future research needs to pay more attention to the effects of transport under cold temperature and loading density on poultry to improve their welfare and reduce economic loss.

Keywords: Transport stress; Poultry; Parameters; Physiological responses; Meat quality

insight into the effects of transport duration, trailer microclimate and loading density on these indicators.

Introduction

The increased rearing number of poultry and the demand for poultry to be centered and slaughtered in processing plants make transport a critical constituent in poultry industry all over the world. Recently there are various concerns about poultry transport due to these three major facts [1,2]. Firstly, mortality may be raised by different transport. Secondly, stressful transport reduces poultry welfare. Thirdly, transport decreases meat quality.

However, transport stressors include feed and water withdrawal, handling, (un)loading, crating density, transport time, ambient temperature, vehicle design, trailer microclimate and lairage time. Among them, transport duration, trailer microclimate and loading density are the three main causes of mortality of birds [3-5].

Stress is regarded to be adaptive or protective responses in body, which protects animals from adverse effects of the stressors. Stress responses, including transport stress responses, are mediated by the two major physiological systems, the Sympatho Adrenal Medullary System (SAMS) and the Hypothalamic-Pituitary-Adrenal Axis (HPA). This can alter the physiological states of animals and subsequently affect meat quality. Transport stress impacts on parameters of bird physiological responses, like corticosterone (CORT), glucose and creatine kinase (CK) before slaughter [6-8] and thus has an effect on meat quality indices such as meat color and water holding capacity (WHC) [9,10]. According to these indicators, we can evaluate the stress level of poultry, so as to improve the management of transport conditions to meet the needs of poultry [6].

Overall this review is aimed to briefly introduce indices of physiological responses and meat quality of birds and to have an

Transport Duration

Transport duration (distance) has an influence on welfare and death loss of birds. Most studies have correlated long-term transport with higher mortalities [3,11-14]. Aral et al. found that for broiler chickens transported for slaughter in Turkey, mortality rates ranged from 0.29% in journeys lasting less than 2 h to 0.46% for journeys lasting more than 10 h [11].

Concentrations of hormones, metabolites and meat color, etc. were also investigated in other studies to evaluate effects of transport time on level of stress and meat quality in birds. Transport stress can activate HPA, increasing the concentrations of blood CORT of broilers [8,15]. Interestingly, a 3 h recovery after transport could facilitate plasma CORT to return to its normal level [16]. Generally, transport stress can decrease lymphocyte quantity and meanwhile increase numbers of heterophils, leading to an elevation on the heterophil:lymphocyte (H:L) ratio [17]. Consequently, the increasing of HL ratio is a reliable and steady parameter reflecting poultry transport stress [6,18,19]. Transport duration could considerably affect plasma glucose level [16,20-22]. Zhang et al. reported that blood glucose increased for short time transport, which was supplied by liver glycogen and a 3 h transport could decrease plasma glucose level, because of the exhaustion of liver glycogen [16]. Similar reductions of glucose following long transport duration were also found by Suchy et al., Sarkar et al. and Vosmerova et al. [20,21,23]. When the breakdown of liver glycogen could not make up for the shortage of glucose, animal body initiates the use of fat and protein. Non-esterified fatty acid (NEFA) is a good indicator of lipolysis. Nijdam et al. reported that blood NEFA level was raised in broilers suffering transport with feed withdrawal compared with those have access to food and also found

that the combination of feed withdrawal and transport consumed more energy of broilers [8]. Besides that plasma uric acid (UA) is a major antioxidant and the primary end product of nitrogen metabolism in poultry, and it is an indicator of oxidative stress and tubular function of the kidney [24]. Huff et al. reported that transport caused a significant increase in turkey plasma UA concentration, indicating a negative energy balance [25]. Creatine kinase (CK) is an intracellular enzyme and its increasing activity is caused by damage to muscle membrane integrity related to stress [7]. The release of CK occurred in ducks following a 2 h transport [26]. Moreover, the increasing concentrations of plasma total protein and albumin are utilized to measure the dehydration of poultry following long time transport. Total protein and albumin levels increased in ducklings due to the 4, 8 and 12 h of the transport [18].

Meat pH and water holding capacity (WHC) are two of the most common parameters for assessing meat quality of poultry. WHC, the ability to keep innate water, is a key property of meat for its influence on meat quality and further processing. In poultry, drip loss is a more frequently used method for WHC. Poultry meat with low pHu (ultimate pH) is related with low WHC, which can alter the color and texture, ultimately affecting consumers' acceptance [27]. Debut et al. stated that if birds were transported for 2 h, the thigh meat showed darker color, high pH and higher curing cooking yield, but transport has no major adverse effect on the breast meat quality [28]. Ducks breast muscle pH 10 min (10 min post-mortem pH) after slaughter was significantly lower after a 2 h transport and reduced muscle quality is also disadvantageous for the duck industry [26]. However, Fernandez et al. gave the results that 30 or 150 min transport duration had little influence on muscle pH 20 min and pH 24 h [29]. The contrasting reports of meat quality could be explained by the not unified measurement methods (different device, measuring time or muscle, etc.) used in experiments and the various transport time, which can influence the results of meat quality after transport.

Furthermore, transport could affect broilers muscle glycogen contents and thus impact on meat quality. A 3 h transport decreased glycogen contents with an increase in lactate content in broilers breast muscle, accompanied with lower pH 24 h and higher drip loss, so transport induced negative changes on broiler breast meat quality could result from the glycolytic enhancement [30]. Savenije et al., Owens et al. and Debut et al. also found the similar results of glycogen contents in birds breast muscle [10,28,31]. However, the increasing transport time decreased the glycogen content of biceps femoris muscle, while that of the breast muscle did not change, possibly since the thigh muscle is involved in maintaining balance in the moving vehicle during transport [32].

Trailer Microclimate

Poultry welfare during transport is most importantly affected by trailer microclimate which includes heat stress (due to high temperature and humidity) and cold stress (due to cold air and wet feathers) and they are the two major reasons for both higher mortality and transport stress [33-36].

A lot of previous work has been done in Europe and thus focuses on heat stress of poultry during transport under medium or high temperature [6,7,15,37,38]. Heat stress during transport increased blood pH and decreased pCO₂ (partial pressure of carbon dioxide) because respiratory over ventilation causes higher emission of carbon dioxide and consequently raises blood pH level [6]. Furthermore, mediated by the stress mechanism, parameters of stress responses and

metabolism, such as CORT, HL, glucose and CK, are varied with the magnitude of the heat load upon the birds [7,15,37,38]. Most research studies have shown acute or seasonal heat stress can result in the declines in pH and WHC and also an increase in lightness of poultry meat [39-41].

In recent years, some researchers noticed transport under extremely cold temperature with poor ventilation inside the trailers. Knezacek et al. reported that at the crate temperature of 17.7 to 55.2°C above than environmental temperature in winter with poor ventilation, broilers experienced both heat stress that occurred in insufficiently ventilated area and cold stress nearby air inlets [42]. However, very little is known about how cold ambient temperature impacts on poultry welfare during transport [43]. Dadgar et al. found that the exposure temperature below 0°C negatively influenced meat quality, such as higher pHu and WHC and darker color, which increased the opportunities of the dark, firm, dry (DFD) meat occurrence [44]. Dadgar et al. also found that 5 and 6 week old broilers were transported at temperatures below -8°C and -14°C during winter, which caused harm effects on meat quality, respectively [45].

Loading Density

High loading density of the trailer could cause heat stress and higher death loss in birds [3,4,46,47]. Mostly during hot climates, more than seven birds per crate can increase the mortality rate [46]. Optimal loading density has an active effect on thermoregulation for transported poultry. However, optimal loading density for poultry transport has not been determined yet, because the effects of loading density are influenced by many factors, for example, the ambient temperature, trailer ventilation, bird size and the position of the drawer within a loaded truck [5,42,47,48]. Lower density means that there are more opportunities for the poultry to regulate body temperatures behaviorally, but too much space per animal may take longer time to load, cost more transport fee, and increase the risk of physical collisions and mechanical bruise [31,49].

Despite of these considerations, there is a little information about effects of different loading densities on physiological responses and meat quality. Delezie et al. reported that transport at high densities resulted in the increasing concentrations of plasma CORT, UA and NEFA, and a decrease of plasma triglyceride concentration, which suggests that the high loading densities caused high stress level and more energy consumption of broilers [15,49,50].

Conclusion

Transport stress is a common issue facing poultry industry today, as it can affect welfare and mortality of birds. This paper summarized the effects of transport duration, trailer microclimate and loading density on physiological responses and meat quality of poultry. Based on this paper, there are two suggestions: (1) The contradictory results of meat quality after transport could be explained by the varied measurement methods and different conditions of application. (2) More future studies need to focus on the effects of loading density and transport under cold temperature on welfare and meat quality of birds.

References

1. Mitchell MA, Kettlewell PJ (2009) Welfare of poultry during transport-a review. Proceedings of 8th Poultry Welfare Symposium, Cervia, Italy.

2. Spurio RS, Soares AL, Carvalho RH, Silveira Junior V, Grespan M, et al. (2016) Improving transport container design to reduce broiler chicken PSE (pale, soft, exudative) meat in Brazil. *Anim Sci J* 87: 277-283.
3. Nijdam E, Arens P, Lambooi E, Decuypere E, Stegeman JA (2004) Factors influencing bruises and mortality of broilers during catching, transport and lairage. *Poult Sci* 83: 1610-1615.
4. Chauvin C, Hillion S, Balaine L, Michel V, Peraste J, et al. (2011) Factors associated with mortality of broilers during transport to slaughterhouse. *Animal* 5: 287-293.
5. Schwartzkopf-Genswein KS, Faucitano L, Dadgar S, Shand P, Gonzalez LA, et al. (2012) Road transport of cattle, swine and poultry in North America and its impact on animal welfare, carcass and meat quality: A review. *Meat Science* 92: 227-243.
6. Mitchell MA, Kettlewell PJ (1998) Physiological stress and welfare of broiler chickens in transit: Solutions not problems! *Poult Sci* 77: 1803-1814.
7. Mitchell MA, Kettlewell PJ, Maxwell MH (1992) Indicators of physiological stress in broiler chickens during road transportation. *Animal Welfare* 1: 91-103.
8. Nijdam E, Delezie E, Lambooi E, Nabuurs MJA, Decuypere E, et al. (2005) Feed withdrawal of broilers before transport changes plasma hormone and metabolite concentrations. *Poult Sci* 84: 1146-1152.
9. Leheska JM, Wulf DM, Maddock RJ (2002) Effects of fasting and transportation on pork quality development and extent of postmortem metabolism. *J Anim Sci* 80: 3194-3202.
10. Owens CM, Sams AR (2000) The influence of transportation on turkey meat quality. *Poult Sci* 79: 1204-1207.
11. Aral Y, Arikian MS, Akin AC, Kuyululu CK, Guloglu SC, et al. (2014) Economic losses due to live weight shrinkage and mortality during the broiler transport. *Ankara Univ Vet Fak Derg* 61: 205-210.
12. Voslarova E, Hytychova T, Vecerek V, Nenadovic K, Bedanova I (2016) Transport-induced mortality in Pekin ducks transported for slaughter in the Czech Republic. *Acta Veterinaria Brno* 85: 205-212.
13. Oba A, Almeida Md, Pinheiro JW, Ida EI, Marchi DF, et al. (2009) The effect of management of transport and lairage conditions on broiler chicken breast meat quality and DOA (Death on Arrival). *Brazilian Archives of Biology and Technology* 52: 205-211.
14. Vieira FMC, Silva IJ, Barbosa Filho JAD, Vieira AMC (2010) Productive losses on broiler preslaughter operations: effects of the distance from farms to abattoirs and of lairage time in a climatized holding area. *Revista Brasileira de Zootecnia* 39: 2471-2476.
15. Freeman BM, Kettlewell PJ, Manning AC, Berry PS (1984) Stress of transportation for broilers. *Vet Rec* 114: 286-287.
16. Zhang L, Yue HY, Zhang HJ, Xu L, Wu SG, et al. (2009) Transport stress in broilers: I. Blood metabolism, glycolytic potential and meat quality. *Poult Sci* 88: 2033-2041.
17. Gross WB, Siegel HS (1983) Evaluation of the heterophil/lymphocyte ratio as a measure of stress in chickens. *Avian Dis* 27: 972-979.
18. Erisir Z, Poyraz O, Erisir M, Onbasilar EE, Erdem E (2008) The changes of the body weight and some blood parameters of Pekin ducklings dependent on transportation duration. *Journal of Animal and Veterinary Advances* 7: 1190-1195.
19. Maxwell MH (1993) Avian blood leucocyte responses to stress. *Worlds Poult Sci J* 49: 34-43.
20. Suchy P, Bedanova I, Vecerek V, Voslarova E, Pistekova V, et al. (2007) Effects of transport stress and floor space reduction on selected biochemical indices in common pheasant (*Phasianus colchicus*). *Archiv fur Geflugelkunde* 71: 56-61.
21. Vosmerova P, Chloupek J, Bedanova I, Chloupek P, Kruzikova K, et al. (2010) Changes in selected biochemical indices related to transport of broilers to slaughterhouse under different ambient temperatures. *Poult Sci* 89: 2719-2725.
22. Yue HY, Zhang L, Wu SG, Xu L, Zhang HJ, et al. (2010) Effects of transport stress on blood metabolism, glycolytic potential and meat quality in meat-type yellow-feathered chickens. *Poult Sci* 89: 413-419.
23. Sarkar P, Ghosh S, Batabyal S, Chatterjee S (2013) Biochemical stress responses of broiler chickens during transport. *Indian J Anim Res* 47: 29-34.
24. Hartman S, Taleb SA, Geng T, Gyenai K, Guan X, et al. (2006) Comparison of plasma uric acid levels in five varieties of the domestic Turkey, *Meleagris gallopavo*. *Poult Sci* 85: 1791-1794.
25. Huff GR, Huff WE, Rath NC, Anthony NB, Nestor KE (2008) Effects of *Escherichia coli* challenge and transport stress on hematology and serum chemistry values of three genetic lines of turkeys. *Poult Sci* 87: 2234-2241.
26. Zhu Z, Chen Y, Huang ZY, Zhang Y, Xu Q, et al. (2014) Effects of transport stress and rest before slaughter on blood parameters and meat quality of ducks. *Canadian Journal of Animal Science* 94: 595-600.
27. Droval AA, Benassi V, Rossa A, Prudencio S, Paião F, et al. (2012) Consumer attitudes and preferences regarding pale, soft, and exudative broiler breast meat. *The Journal of Applied Poultry Research* 21: 502-507.
28. Debut M, Berri C, Baeza E, Sellier N, Arnould C, et al. (2003) Variation of chicken technological meat quality in relation to genotype and preslaughter stress conditions. *Poult Sci* 82: 1829-1838.
29. Fernandez X, Bouillier-Oudot M, Molette C, Bernadet MD, Manse H (2011) Duration of transport and holding in lairage at constant postprandial delay to slaughter—Effects on fatty liver and breast muscle quality in mule ducks. *Poult Sci* 90: 2360-2369.
30. Zhang L, Li JL, Gao T, Lin M, Wang XF, et al. (2014) Effects of dietary supplementation with creatine monohydrate during the finishing period on growth performance, carcass traits, meat quality and muscle glycolytic potential of broilers subjected to transport stress. *Animal* 8: 1955-1962.
31. Savenije B, Lambooi E, Gerritzen MA, Venema K, Korf J (2002) Effects of feed deprivation and transport on preslaughter blood metabolites, early postmortem muscle metabolites and meat quality. *Poult Sci* 81: 699-708.
32. Warriss PD, Kestin SC, Brown SN, Knowles TG, Wilkins LJ, et al. (1993) The depletion of glycogen stores and indices of dehydration in transported broilers. *Br Vet J* 149: 391-398.
33. Vieira FM, Silva IJ, Barbosa Filho JA, Vieira AM, Broom DM (2011) Preslaughter mortality of broilers in relation to lairage and season in a subtropical climate. *Poult Sci* 90: 2127-2133.
34. Hunter R, Mitchell M, Carlisle A (1999) Wetting of broilers during cold weather transport: A major source of physiological stress? *Br Poult Sci* 40: 48-49.
35. Hunter R, Mitchell M, Matheu C (2001) Mortality of broiler chickens in transit—correlation with the thermal micro-environment. *Livestock Environment VI Proc 6th Int Symp. Am Soc Agric Eng Louisville: Kentucky*.
36. Vecerek V, Grbalova S, Voslarova E, Janackova B, Malena M (2006) Effects of travel distance and the season of the year on death rates of broilers transported to poultry processing plants. *Poult Sci* 85: 1881-1884.
37. Mitchell M, Kettlewell P, Hunter R, Carlisle A (2001) Physiological stress response modeling—application to the broiler transport thermal environment. *International Symposium*.
38. González V, Rojas G, Aguilera A, Flores-Peinado S, Lemus-Flores C, et al. (2007) Effect of heat stress during transport and rest before slaughter, on the metabolic profile, blood gases and meat quality of quail. *Int J Poult Sci* 6: 397-402.
39. Holm CP, Fletcher DL (1997) Antemortem holding temperatures and broiler breast meat quality. *J Appl Poult Res* 6: 180-184.
40. Petracci M, Fletcher DL, Northcutt JK (2001) The effect of holding temperature on live shrink, processing yield and breast meat quality of broiler chickens. *Poult Sci* 80: 670-675.
41. Mitchell M (2006) Influence of pre-slaughter stress on animal welfare and processing efficiency. *Worlds Poult Sci J* 62: 254.
42. Knezacek T, Olkowski A, Kettlewell P, Mitchell M, Classen H (2010) Temperature gradients in trailers and changes in broiler rectal and core body temperature during winter transportation in Saskatchewan. *Can J Anim Sci* 90: 321.

-
43. Strawford ML, Watts JM, Crowe TG, Classen HL, Shand PJ (2011) The effect of simulated cold weather transport on core body temperature and behavior of broilers. *Poult Sci* 90: 2415-2424.
 44. Dadgar S, Lee ES, Leer TLV, Burlingette N, Classen HL, et al. (2010) Effect of microclimate temperature during transportation of broiler chickens on quality of the pectoralis major muscle. *Poult Sci* 89: 1033-1041.
 45. Dadgar S, Lee ES, Leer TL, Crowe TG, Classen HL, et al. (2011) Effect of acute cold exposure, age, sex and lairage on broiler breast meat quality. *Poult Sci* 90: 444-457.
 46. Vieira FMC, Silva IJO, Barbosa Filho JAD, Vieira AMC (2013) Reducing poultry preslaughter losses: Crating density effects under different lairage periods at slaughterhouse. *JABB-Online Submission System* 1: 1-6.
 47. Whiting TL, Drain ME, Rasali DP (2007) Warm weather transport of broiler chickens in Manitoba. II. Truck management factors associated with death loss in transit to slaughter. *Can Vet J* 48: 148.
 48. Yalçın S, Özkan S, Oktay G, Çabuk M, Erbayraktar Z, et al. (2004) Age-related effects of catching, crating and transportation at different seasons on core body temperature and physiological blood parameters in broilers. *J Appl Poult Res* 13: 549-560.
 49. Delezie E, Swennen Q, Buyse J, Decuypere E (2007) The effect of feed withdrawal and crating density in transit on metabolism and meat quality of broilers at slaughter weight. *Poult Sci* 86: 1414-1423.
 50. Siegel Hv, Van Kampen M (1984) Energy relationships in growing chickens given daily injections of corticosterone. *Br Poult Sci* 25: 477-485.