

Serotonin 4 Receptors: A Cornerstone in Anorexia Nervosa?

Valérie Compan*

Nîmes University, France

Abstract

Adaptive decision-making to eat is crucial for survival but in anorexia nervosa, the brain persistently supports reduced food intake despite the physiological need to consume food. How the brain persists in reducing food intake sometimes even to the point of death despite the evolution of multiple mechanisms to ensure survival by governing adaptive eating behaviors remains mysterious. Food intake is a conserved behavioral trait between all species and involves numerous biological systems including the old phylogenetically serotonergic system. The present review focuses on anorexia and the implication of specific serotonin (5-HT, 5-hydroxytryptamine) receptors in food intake. In this context, we found that an early restrictive food intake due to stress, critically engages goal-directed (decision-making) systems upon the control of the serotonin 5-HT₄ receptors, supporting that an early food restriction may first protect from depressive-like states but could become a deadly dependence. Finally, in the face to environmental challenges, an initial protective and beneficial adaptive response could become a pathologic dependence.

Keywords: Behavioral trait; Anorexia nervosa; Depression

Introduction

How does the brain implement inappropriate eating decisions to the point of starvation and death even though it has evolved to favor adapted and adaptive eating behaviors for survival? Solving this mystery poses a vital challenge because restrictive feeding aggravates numerous diseases (e.g. cancer, diabetes type I) and places its extreme aspect, anorexia nervosa, among the first cause of death of adolescents in Europe [1]. As for most behavioral impairments, the cause may depend on a crosstalk between environmental factors and a biological predisposition.

In the face of environmental changes, behavioral disturbances are often correlated to deregulations of neural circuits and explored in animals' brain, permitting to study phenotypes in isolation from molecular to behavioral traits. Human mutations exist in mental diseases and the corresponding mouse models reveal the conservation of specific mechanisms because some treatments are efficient to reduce some behavioral traits [2-4]. These commonalities could rely on old phylogenetically biological systems.

Food intake is a conserved behavioral trait between all species and involves numerous biological systems including one of the most conserved neural systems, the serotonergic system, known for controlling particular aspects of feeding behavior in both rodents and humans. In mammals, the serotonergic neuronal cell bodies and dendrites assemble in the raphe nuclei. Among nine nuclei, the dorsal and median raphe nuclei (DR, MR) send axons to the whole forebrain [5]. In particular, the serotonergic axons in the cerebral cortex mainly arise from the DR [6]. Serotonin (5-hydroxytryptamine: 5-HT) binds 18 G-protein coupled receptor subtypes (5-HTRs), more often located at 100 μm [7] than at 20 nm (synaptic transmission) up to from the site of 5-HT release. The preponderant 5-HT volume transmission extends the ubiquitous distribution of the 5-HT system and likely their plural functional implications. For example, this neural system is critical for regulating molecular substrates of survival, i.e., preventing depressive-like states, anxiety-like behaviour, i.e., the fear to novelty, locomotion, learning and memory, including peripheral functions such as the gastro-intestinal peristalsis and food intake. Deregulation of the 5-HT systems appear then as a critical support of the different symptoms seen in patients with anorexia nervosa.

The present review argues that the neural substrates of an early anorexia-like behavior could limit depressive-like states, i.e., favoring the activity of protective mechanisms of survival upon the control of the serotonin 4 receptors (5-HT₄Rs). Chronic external stress could however challenge the limits of the neuronal plasticity and favor an addiction (without drugs; a dependence) to anorexia. A first part summarizes the symptomatology of anorexia nervosa, as previously described in detail elsewhere [8]. External stress often precedes the occurrence of anorexia nervosa, as well as other diseases. Consequently, our scientific community has used stressed-animal models with anorexia-like behavior, as summarized in a second part. A third part then summarizes the first example of causal relationships between specific molecular events and a reduced food intake due to stress. These neural bases appear to mainly depend on the activity of the efferent 5-HT neurons from the dorsal raphe nucleus (DR) to the ventral medial prefrontal cortex (mPFC), a crucial pathway controlling adaptive responses to stress. In this context and among numerous factors, the serotonin 4 receptors (5-HT₄Rs) represent, in animals, a causal link between an early anorexia-like behavior and stress and, a potential therapeutic target of this actual deadly disease.

Symptomatology of Anorexia Nervosa

Patients with anorexia nervosa can reduce food intake until death, display emaciation, amenorrhea, motor hyperactivity or "over-exercise" [9,10] express anxiety [11,12] harm avoidance [13], perfectionism [14], obsession [15,16] and often suffer from depression [17]. Individuals suffering from anorexia nervosa can also struggle with bulimia, i.e., overeating with purging. The symptomatology of anorexia nervosa is extremely complex with a critical absence of medication [18] and different medical complication, including substance abuse and suicide

*Corresponding author: Valérie Compan, Nîmes University, Place Gabriel Peri, Carmes, 30021 Nîmes, France, Tel: 33664915140; E-mail: valerie.compan@unimes.fr

Received March 31, 2017; Accepted May 09, 2017; Published May 16, 2017

Citation: Compan V (2017) Serotonin 4 Receptors: A Cornerstone in Anorexia Nervosa? Autism Open Access 7: 207. doi:10.4172/2165-7890.1000207

Copyright: © 2017 Compan V, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

[19,20]. Food restriction can even alternate with bulimia that differs from binge eating [21]. Binge eating involves uncontrollable consumption of large amounts of food but is not followed by food purging. Personality traits (anxiety, harm avoidance, obsession, perfectionism), occurring in childhood before the onset of eating disorders, likely reveal a progressive implementation of a biological predisposition, which could, when associated to genetic heritability, account for approximately 50-80% of the risk of developing eating disorders [22,23] when facing environmental changes (stress). The symptomatology of anorexia nervosa is then extremely complex. The term of "anorexia" is often used instead of "anorexia nervosa", likely because "anorexia" is the hallmark symptom. We also use "anorexia" instead of "anorexia-like behavior" for animals. Animals display anorexia that is operationally defined as reduced food intake despite the physiological energy demand, i.e., following partial or total food deprivation [24]. These animal models do not recapitulate all the symptoms of anorexia nervosa but few of them, as described below. However, this is noteworthy to see how the implication of the 5-HT_{2c}R [25,26] and the MCR4 receptors [27-30] is conserved from rodents to humans, suggesting commonalities that could rely on phylogenetically old neural systems. These studies highlight that the importance to study phenotypes in isolation from molecular to behavioural traits using animal models in order to propose some potential effective therapeutic target in humans.

Intercross between a Biological Predisposition and Environmental Challenges

Food intake in both animals and humans are stress-dependent [31,32]. One of the most employed animal models to identify the neural basis of a reduced food intake in fed ad libitum rodents, i.e., hypophagia, due to stress is forced immobilization, called the restraint stress [31,33-45]. The ability of stress to trigger reduction in food intake has been attributed to an increase in the activity of the serotonergic systems but also to the hyperactivity of hypothalamo-pituitary adrenal axis. Peptides of the corticotropin-releasing hormone (CRH) family, such as stresscopin or urocortin, induce decreases in food intake. The reciprocal influences between serotonergic systems and the stress axis have made it difficult to identify a clear neurochemical cascade underlying the influence of stress on feeding behavior. A working hypothesis would be that an increase in the activity of the HPA axis could induce an elevation in 5-HT, which in combination with stress hormones induce a decrease in food intake. In keeping with this hypothesis CRH has been shown to stimulate the activity of serotonergic neurons. In addition, repeated injections of corticosterone enhance the excitatory effect of agonists of the 5-HT₄Rs on hippocampal CA1 neurons. In humans, changes in the levels of cortisol and the concentration of the 5-HT₄Rs appear negatively correlated [46]. Increased levels of cortisol (50%) correlate a reduction in the concentration of the 5-HT₄Rs in the prefrontal and anterior cingulate cortex, pallidostriatum, 20-30 min following awakening in the morning. Such a reduction in the concentration of the 5-HT₄Rs could succeed an elevation in the levels of 5-HT in the synaptic cleft because a 5-HT depletion enhances the levels of the 5-HT₄Rs in different brain areas (basal ganglia) in both rodents and in humans [47,48]. How the 5-HT₄Rs may intervene in the regulation of cortisol (and conversely) remains to be fully investigated. A correlation between modifications in the levels of 5-HT, the 5-HT₄Rs and cortisol appears to exist and be critical for regulating food intake following stress but the exact mechanisms remain undetermined.

In this context, despite the ability of CRH to reduce food intake, mice lacking the CRH remain sensitive to stress-induced hypophagia [38]. The 5-HT₄R knockout (KO) mice did not display a general maladaptive

response to stress because changes in the levels of corticosterone (and glucose) following stress are similarly increased compared with wild-type (WT) animals [42]. One could then focus on the serotonergic system in order to better identify the neural basis of hypophagia due to stress.

Numerous studies relate an increase in 5-HT volume transmission and stress -induced anorexia-like behavior. Stress-related behavioral paradigms such as conditioned fear increase 5-HT metabolism and release in the mPFC, nucleus accumbens (NAc), amygdala and dorsal hippocampus. In particular, the restraint stress increases the 5-HT turnover in the hypothalamus and amygdala in mice and rats. Only two 5-HT receptors appear critically involve in stress-induced anorexia-like behavior because mice treated with 8-OH-DPAT [34], a 5-HT_{1A}R/5-HT₇R agonist or lacking the 5-HT₄Rs exhibit an attenuated hypophagia [42].

Serotonin Volume Transmission Reduced Food Intake

The 5-HT system commonly mediates reduction in food intake, i.e., hypophagia [8,49]. Stimulating the Gi-coupled 5-HT_{1A} receptors in the DR (DR-5-HT_{1A}R) reduces the firing activity of DR 5-HT neurons, and mediates hyperphagia [49]. Most of studies, conducted in the hypothalamus, describe animals with hypophagia following stimulation of the 5-HT_{1B} and 5-HT_{2C} receptors [5-HT_{1B}R, 5-HT_{2C}R] [8] while, stimulation of the 5-HT_{1A}R and 5-HT_{2B}R can exceptionally induce hyperphagia [50]. The serotonergic system can also hijack a motivation for food in food-deprived mice, mimicking an anorexia-like behavior. This involves the activation of an addictive signaling pathway [cAMP-protein kinase A (PKA)/CART: cocaine- and amphetamine-regulated transcript] upon the control of the 5-HT₄Rs in the NAc, a critical structure in the brain's reward system [24,51]. In addition, mice lacking 5-HT_{1B}R self-imposed food restriction after deprivation and still displayed anorexia-like behavior and hyperactivity after MDMA (3,4-N-methylenedioxymethamphetamine, the psychogenic compound of ecstasy [24]. Decryption of the mechanisms showed a gain-of-function of the 5-HT₄Rs in the absence of 5-HT_{1B}R, associated with CART overexpression in the NAc and not in other brain areas [24]. The implication or not of the different 5-HTRs in food intake also depends on their localization in the brain [52]. It can be further complex because alternative splicing of 5-HT_{2C}R induced by a systemic delivery of oligonucleotide can reduce food intake [53].

In humans, the inhibitory control of food intake through the 5-HT volume transmission is retained [8,49,54]. In particular, patients who had recovered from one of the symptoms of anorexia nervosa, i.e., from a persistent food restriction (called anorexia), display an increase in the activity of the DR-5-HT_{1A}R [55]. Obesity related to hyperphagia correlates 5-HT depletion and compensatory increases in the levels of NAc-5-HT₄Rs in both rats and humans [47-49,56,57].

The Serotonin 4 Receptors: A Promising Therapeutic Target

Food intake clearly depends on the activity of the 5-HT system and, both food intake and activity of the 5-HT system depend on external stressors [49,58]. However, whether changes in food intake and the activity of the 5-HT system in the face to external stress are causally related or correlated, remain undetermined, with the exception of a possible network controlled by the 5-HT₄Rs, as proposed here. Following up a summary of unpublished results yet, one may suspect that the neural network of an "early anorexia" [59] in the face to external stress can protect the brain from depressive-like states.

The cerebral distribution of the 5-HT₄Rs is conserved in humans, with the highest levels in the shell part of the NAc and the lowest, in the cerebral cortex [47,60-62]. The 5-HT₄Rs exert a positive feedback on the DR-5-HT cells, not from the DR (they are absent) but from the mPFC. The first example of functional implication of the 5-HT₄Rs in brain has been first described in the 5-HT₄R knockout (KO) mice [42], with the exception of a primary description of the positive implication of the 5-HT₄Rs on associative memory [63].

The 5-HT₄R KO mice better resist stress-induced reduction in food intake, i.e., hypophagia [42]. The attenuated hypophagia in stressed 5-HT₄R KO mice was accompanied by a reduced motor reactivity to novelty [42]. Consistently, stimulating (or overexpressing) the 5-HT₄Rs in the NAc reduces food intake and increases motor hyperactivity [24]. As mentioned above and recall here, anorexia-like behavior and motor hyperactivity following stimulation of the 5-HT₄Rs in the NAc depends on the activation of an addictive signaling pathway (cAMP/PKA/CART) [24,51]. The cAMP/PKA signaling pathway is critical in cocaine addiction [64]. As we described elsewhere in detail [8] and recall here, we have detected common molecular mechanisms between anorexia and addiction. Indeed, drugs of abuse (e.g. cocaine, amphetamine) trigger adaptive responses including an increased activity of the cAMP/PKA signaling pathway in the NAc [65-67]. The resultant phosphorylation of the cAMP-responsive element binding (CREB) dampens rewarding effects. Consequently, the sensitivity to subsequent drug exposures decreases (tolerance) with increased activity of reward pathways (dependence) to the point that drugs removal triggers motivation decline, mimicking depressive states [68]. Stimulation of the cAMP/PKA/CART pathway, in the NAc, following local stimulation of the 5-HT₄R provokes anorexia [51]. This pathway is also involved in anorexia induced by the 3,4-N-methylenedioxymethamphetamine (MDMA), the psychogenic compound of ecstasy. The ability of cocaine addiction-related animal models [69] to self-impose food restriction despite an early period of deprivation further depends on a gain-of-function of 5-HT₄Rs with CART overexpression in the NAc [24]. Considering the involvement of CART in motivational properties of cocaine [70,71], these findings evidence a “shared neural signal foul-up” between drug dependence and anorexia, consistent with deficits in neural networks underlying addiction in patients with anorexia nervosa [23,72]. Interestingly, anorexia and motor hyperactivity are two hallmarks of anorexia nervosa, suggesting a prime dysfunction of the 5-HT₄R in anorexia nervosa. Male mice with a 5-HT₄R overexpression, in the NAc, persist in self-restricting despite an earlier imposed period of partial (-20%) food deprivation [24].

Consistently with the ability of the 5-HT₄Rs to activate an addictive (a dependent) signaling pathway (cAMP/PKA/pCREB: cAMP-responsive element binding protein phosphorylation) in the NAc, mice lacking the 5-HT₄Rs display an anhedonia-like behavior [73]. CREB overexpression in the NAc reduces the rewarding effects of sucrose [74] and the ability of cocaine to induce CREB phosphorylation is absent in the NAc of the 5-HT₄R KO mice [8]. The absence of 5-HT₄Rs could then favor an anhedonic behavior while their hyperactivity could mediate a rewarding effect. Steinglass et al. has suspected an initial rewarding effect of an early anorexia, helping patients to manage stressors [75], consistently with other clinical studies [23,76].

Many drugs of abuse also induce anxiety-like behavior. It is appropriate, here, to mention both a high level of anxiety in patients with anorexia nervosa [77] and the critical implication of 5-HT systems in anxiety-like behavior. This emotional disturbance is prevented when the 5-HT_{1A}R is especially expressed during the early postnatal

period [78]. In contrast, the absence of 5-HT₄Rs induced anxiety-like behavior in stressful conditions and leads to decrease in the levels of 5-HT_{1A}R in the dorsal hippocampus [42,79]. Interestingly, the 5-HT₄R clearly enhances pCREB in the NAc (Compan, results). Considering that inhibition of the transcription factor CREB in the NAc has been associated with anxiety-like behavior, anorexia induced by stimulation of 5-HT₄R in the NAc could favor the “anxiety-reducing character to dietary restraint” [80-82]. In contrast, reduced activation of 5-HT₄R could enhance anxiety that is provoked by overeating [8].

A key question is what are the neural substrates that initiate the transition from transient to persistent restrictive food intake, i.e., those that would abnormally favor an “early decision-making” to not eat. We suspected impaired activity of a network governing goal-directed behavior (decision-making), i.e., the ascending serotonergic inputs from the DR to the mPFC for several reasons: (i) this network supports adaptive “decision” to avoid adverse effect of stress and then depressive states [83-85] (ii) eating disorders are stress-related [58] (iii) and often associated with depression [86]. Our working hypotheses and our most recent results are schematized in the enclosed video [87], introducing that processes within the voluntary nervous system (underlying decision, motivation) could be modified to prevail over a cerebral autonomous control of hunger, compromising survival.

Results include an abnormal absence of reduction in the 5-HT₄Rs in the NAc in stressed animals when dispossessed of the 5-HT₄Rs only in the mPFC (Compan, results), suggesting that an abnormal overexpression could occur in the NAc and favor a dependent anorexia. This study suggests a primary mechanism that could support the onset of a persistent hypophagia [“an early anorexia”], whereby individuals shift from adaptive to persistent maladaptive food choice [88].

Therefore, examination of the activity of neural centers involved in the recognition of rewards and the development of habits is relevant [88]. Accordingly, food choices to the point of starvation correlates with feeding habits and engaged the dorsal striatum [89]. A recent report described goal-directed decision-making as a complex process and argued that reward-based decisions depend on the habit and goal-oriented systems [90]. The habit system “stores” stimulus-response associations based on past rewards and the goal-oriented system selects one action by anticipating the positive and negative outcomes [90]. Since additionally, “addiction is a form of learning and relapse is a persistent memory of the drug experience” [91]. Interestingly, the implication of the 5-HT₄R in memory is well documented, as summarized elsewhere [92] are implicated in long-term memory [93]. The 5-HT₄R KO mice performed similarly for spatial learning, short- and long-term retention related to adaptive changes in cholinergic systems but the loss-of-function of the 5-HT₄Rs aggravates scopolamine-induced deficits in long-term memory [93]. Interestingly, a recent study suggests a positive influence of the 5-HT₄R in the NAc on emotional memory consolidation [94].

Finally, this is important to note, here, some medical complications in patients with anorexia nervosa related to the gastrointestinal tract. Again, the 5-HT₄Rs contribute to control peripheral effects on the gastro-intestinal tract where they may serve as targets for treatment of dyspepsia, gastro-oesophageal reflux disease, gastroparesis or irritable bowel syndrome [95]. The medication of these peripheral disorders also target the 5-HT₃R, inducing side effects and, it remains unclear whether targeting only the 5-HT₄R would or not be the most relevant strategy [96]. Numerous recent studies focus on the beneficial effect of prucalopride, a partial agonist of the 5-HT₄Rs [97].

Conclusion

Feeding behavior appears to result from an integrated activity of the autonomic and voluntary nervous systems informed by the sensory nervous system of external environment states, which may develop and persist over time. Studying neural substrates of feeding behavior is therefore crucial to better understand how the main nervous systems interact to adapt decision to eat in the face of external environment. In this context, the symptomatology of anorexia nervosa reveals a “crosstalk” between different cerebral structures where the 5-HT₄Rs appear to systematically intervene (rewarding effect of anorexia, motor hyperactivity, anxiety, memory). Mainly based on recent results, the voluntary control processes in the nervous system (underlying decision, motivation) could be modified to prevail over cerebral autonomous control of hunger, compromising survival. We have identified that the neural adaptive responses to stress, known to reduce defective behavior of self-preservation (depression), initiate a persistent hypophagia following stress (video) depend on the 5-HT₄Rs in the mPFC, a critical structure implicated in “decision-making”. An “early anorexia” could then first protect self-preservation via neural pathways concerned with dealing with stress because in itself an overexpression of the 5-HT₄R in the mPFC in stressed animals triggers a part of the molecular effects of an antidepressant and, triggers hypophagia. In the face of chronic stress, limits of this adaptive process could “submerge” cortical control and “release the influence of the subcortical areas” such as the NAc (autonomous control without adaptive decisional control), in which uncontrolled oscillating changes in common molecule levels (cAMP, CREB: all controlled by GPCRs) could lead to a maladaptive consumption of foods. Our most recent study introduces a primary mechanism whereby individuals could shift from transient to persistent food restriction as seen in anorexia nervosa [88] and in agreement with numerous studies make further conceivable to target 5-HT₄Rs to treat this incurable disease. We now can look forward to understanding how the influence of the 5-HT₄Rs in long-term memory [93] - related to their positive control of the cAMP signaling from where come change in neuron morphology (increased number of dendritic spines) - could underlie some behavioral traits in anorexia nervosa. Indeed, nerve growth and synapse formation (induced by pCREB) represent a dynamic process whereby individuals store information for a long period of time (long term memory). Could actions of the 5-HT₄Rs be related to changes in the habit neural system seen in anorexia nervosa [89]? Considering the importance of molecules involved in the developmental formation of synapses and autism (neurexin, neuroligin), could the 5-HT₄Rs underlie rigid attitudes and altered social interactions seen in both symptomatology of autism and anorexia nervosa?

Acknowledgement

The Agence Nationale de la Recherche (ANR-09-MNPS-024-01: SERFEED) and Anorexia, Dependence, Obesity, Receptors Foundation has supported this work.

References

- Papadopoulos FC, Pantziaras I, Lagiou P, Brandt L, Ekselius L, et al. (2009) Age at onset of anorexia nervosa and breast cancer risk. *Eur J Cancer Prev* 18: 207-211.
- Bevilacqua L, Doly S, Kaprio J, Yuan Q, Tikkanen R, et al. (2010) A population-specific HTR2B stop codon predisposes to severe impulsivity. *Nature* 468: 1061-1066.
- Kirov G, Pocklington AJ, Holmans P, Ivanov D, Ikeda M, et al. (2012) De novo CNV analysis implicates specific abnormalities of postsynaptic signalling complexes in the pathogenesis of schizophrenia. *Mol Psychiatry* 17: 142-153.
- Donaldson ZR, Hen R (2015) From psychiatric disorders to animal models: A bidirectional and dimensional approach. *Biol Psychiatry* 77: 15-21.
- Azmitia EC (1999) Serotonin neurons, neuroplasticity and homeostasis of neural tissue. *Neuropsychopharmacology* 21: 33S-45S.
- Muzerelle A, Scotto-Lomassese S, Bernard JF, Soiza-Reilly M, Gaspar P (2016) Conditional anterograde tracing reveals distinct targeting of individual serotonin cell groups (B5-B9) to the forebrain and brainstem. *Brain Struct Funct* 221: 535-561.
- Descarries L, Beaudet A, Watkins KC (1975) Serotonin nerve terminals in adult rat neocortex. *Brain Res* 100: 563-588.
- Compan V, Walsh BT, Kaye W, Geliebter A (2015) How does the brain implement adaptive decision making to eat? *J Neurosci* 35: 13868-13878.
- Beumont PJ, Arthur B, Russell JD, Touyz SW (1994) Excessive physical activity in dieting disorder patients: Proposals for a supervised exercise program. *Int J Eat Disord* 15: 21-36.
- Casper RC (2006) The 'drive for activity' and "restlessness" in anorexia nervosa: Potential pathways. *J Affect Disord* 92: 99-107.
- Godart NT, Flament MF, Lecrubier Y, Jeammet P (2000) Anxiety disorders in anorexia nervosa and bulimia nervosa: Co-morbidity and chronology of appearance. *Eur Psychiatry* 15: 38-45.
- Kaye WH, Bulik CM, Thornton L, Barbarich N, Masters K (2004) Comorbidity of anxiety disorders with anorexia and bulimia nervosa. *Am J Psychiatry* 161: 2215-2221.
- Fassino S, Abbate-Daga G, Amianto F, Leombruni P, Boggio S, et al. (2002) Temperament and character profile of eating disorders: A controlled study with the temperament and character inventory. *Int J Eat Disord* 32: 412-425.
- Friederich HC, Herzog W (2011) Cognitive-behavioral flexibility in anorexia nervosa. *Curr Top Behav Neurosci* 6: 111-123.
- Anderlueh MB, Tchanturia K, Rabe-Hesketh S, Treasure J (2003) Childhood obsessive-compulsive personality traits in adult women with eating disorders: Defining a broader eating disorder phenotype. *Am J Psychiatry* 160: 242-247.
- van den Heuvel OA, Veltman DJ, Groenewegen HJ, Witter MP, Merckelbach J, et al. (2005) Disorder-specific neuroanatomical correlates of attentional bias in obsessive-compulsive disorder, panic disorder and hypochondriasis. *Arch Gen Psychiatry* 62: 922-933.
- Casper RC (1998) Depression and eating disorders. *Depress Anxiety* 1: 96-104.
- Kaye WH, Wierenga CE, Knatz S, Liang J, Boutelle K, et al. (2015) Temperament-based treatment for anorexia nervosa. *Eur Eat Disord Rev* 23: 12-18.
- Comerci GD (1990) Medical complications of anorexia nervosa and bulimia nervosa. *Med Clin North Am* 74: 1293-1310.
- Westmoreland P, Krantz MJ, Mehler PS (2016) Medical complications of anorexia nervosa and bulimia. *Am J Med* 129: 30-37.
- Corwin RL, Avena NM, Boggiano MM (2011) Feeding and reward: Perspectives from three rat models of binge eating. *Physiol Behav* 104: 87-97.
- Bulik CM, Sullivan PF, Tozzi F, Furberg H, Lichtenstein P, et al. (2006) Prevalence, heritability and prospective risk factors for anorexia nervosa. *Arch Gen Psychiatry* 63: 305-312.
- Kaye WH, Fudge JL, Paulus M (2009) New insights into symptoms and neurocircuit function of anorexia nervosa. *Nat Rev Neurosci* 10: 573-584.
- Jean A, Laurent L, Bockaert J, Charnay Y, Dusticier N, et al. (2012a) The nucleus accumbens 5-HTR(4)-CART pathway ties anorexia to hyperactivity. *Transl Psychiatry* 2: e203.
- Higgins GA, Sellers EM, Fletcher PJ (2013) From obesity to substance abuse: Therapeutic opportunities for 5-HT_{2C} receptor agonists. *Trends Pharmacol Sci* 34: 560-570.
- Howell LL, Cunningham KA (2015) Serotonin 5-HT₂ receptor interactions with dopamine function: Implications for therapeutics in cocaine use disorder. *Pharmacol Rev* 67: 176-197.
- Adan RA, Kas MJ (2003) Inverse agonism gains weight. *Trends Pharmacol Sci* 24: 315-321.
- Lubrano-Berthelot C, Cavazos M, Dubern B, Shapiro A, Stunff CL, et al. (2003) Molecular genetics of human obesity-associated MC4R mutations. *Ann N Y Acad Sci* 994: 49-57.

29. Srinivasan S, Lubrano-Berthelie C, Govaerts C, Picard F, Santiago P, et al. (2004) Constitutive activity of the melanocortin-4 receptor is maintained by its N-terminal domain and plays a role in energy homeostasis in humans. *J Clin Invest* 114: 1158-1164.
30. Kim DH, Shin SW, Baik JH (2008) Role of third intracellular loop of the melanocortin 4 receptor in the regulation of constitutive activity. *Biochem Biophys Res Commun* 365: 439-445.
31. Laurent L, Jean A, Manrique C, Najimi M, Chigr F, et al. (2012) Anorexia and drugs of abuse abnormally suppress appetite, the result of a shared molecular signal foul-up. In: *Animal model of eating disorders* 74: 319-331.
32. Hardaway JA, Crowley NA, Bulik CM, Kash TL (2015a) Integrated circuits and molecular components for stress and feeding: implications for eating disorders. *Genes Brain Behav* 14: 85-97.
33. Krahn DD, Gosnell BA, Grace M, Levine AS (1986) CRF antagonist partially reverses CRF and stress-induced effects on feeding. *Brain Res Bull* 17: 285-289.
34. Dourish CT, Kennett GA, Curzon G (1987) The 5-HT_{1A} agonists 8-OH-DPAT, buspirone and ipsapirone attenuate stress-induced anorexia in rats. *J Psychopharmacol* 1: 23-30.
35. Shimizu N, Oomura Y, Kai Y (1989) Stress-induced anorexia in rats mediated by serotonergic mechanisms in the hypothalamus. *Physiol Behav* 46: 835-841.
36. Grignaschi G, Mantelli B, Samanin R (1993) The hypophagic effect of restraint stress in rats can be mediated by 5-HT₂ receptors in the paraventricular nucleus of the hypothalamus. *Neurosci Lett* 152: 103-106.
37. Haleem DJ, Jabeen B, Parveen T (1998) Inhibition of restraint-induced anorexia by injected tryptophan. *Life Sci* 63: PL205-212.
38. Weninger SC, Muglia LJ, Jacobson L, Majzoub JA (1999) CRH-deficient mice have a normal anorectic response to chronic stress. *Regul Pept* 84: 69-74.
39. Wang SW (2002) Effects of restraint stress and serotonin on macronutrient selection: A rat model of stress-induced anorexia. *Eat Weight Disord* 7: 23-31.
40. Chaki S, Ogawa S, Toda Y, Funakoshi T, Okuyama S (2003) Involvement of the melanocortin MC4 receptor in stress-related behavior in rodents. *Eur J Pharmacol* 474: 95-101.
41. Chou-Green JM, Holscher TD, Dallman MF, Akana SF (2003) Repeated stress in young and old 5-HT_{2C} receptor knockout mice. *Physiol Behav* 79: 217-226.
42. Compan V, Zhou M, Grailhe R, Gazzara RA, Martin R, et al. (2004) Attenuated response to stress and novelty and hypersensitivity to seizures in 5-HT₄ receptor knock-out mice. *J Neurosci* 24: 412-419.
43. Kinzig KP, Hargrave SL, Honors MA (2008) Binge-type eating attenuates corticosterone and hypophagic responses to restraint stress. *Physiol Behav* 95: 108-113.
44. Bazhan NM, Yakovleva TV, Kazantseva AY, Makarova EN (2013) Exaggerated anorexigenic response to restraint stress in A(y) mice is associated with elevated CRFR2 mRNA expression in the hypothalamus. *Physiol Behav* 120: 19-25.
45. Haque Z, Akbar N, Yasmin F, Haleem MA, Haleem DJ (2013) Inhibition of immobilization stress-induced anorexia, behavioral deficits and plasma corticosterone secretion by injected leptin in rats. *Stress* 16: 353-362.
46. Jakobsen GR, Fisher PM, Dyssegaard A, McMahon B, Holst KK, et al. (2016) Brain serotonin 4 receptor binding is associated with the cortisol awakening response. *Psychoneuroendocrinology* 67: 124-132.
47. Compan V, Daszuta A, Salin P, Sebben M, Bockaert J, et al. (1996) Lesion study of the distribution of serotonin 5-HT₄ receptors in rat basal ganglia and hippocampus. *Eur J Neurosci* 8: 2591-2598.
48. Haahr ME, Fisher PM, Jensen CG, Frokjaer VG, Mahon BM, et al. (2014) Central 5-HT₄ receptor binding as biomarker of serotonergic tonus in humans: A [¹¹C] SB207145 PET study. *Mol Psychiatry* 19: 427-432.
49. Compan V (2013) Under- to over-eating: What role do serotonin receptors play? *Future Neurology*.
50. Yadav VK, Oury F, Suda N, Liu ZW, Gao XB, et al. (2009) A serotonin-dependent mechanism explains the leptin regulation of bone mass, appetite and energy expenditure. *Cell* 138: 976-989.
51. Jean A, Conductier G, Manrique C, Bouras C, Berta P, et al. (2007) Anorexia induced by activation of serotonin 5-HT₄ receptors is mediated by increases in CART in the nucleus accumbens. *Proc Natl Acad Sci U S A* 104: 16335-16340.
52. Pratt WE, Clissold KA, Lin P, Cain AE, Ciesinski AF, et al. (2016) A systematic investigation of the differential roles for ventral tegmentum serotonin 1 and 2 type receptors on food intake in the rat. *Brain Res* 1648: 54-68.
53. Zhang Z, Shen M, Gresch PJ, Ghamari-Langroudi M, Rabchevsky AG, et al. (2016) Oligonucleotide-induced alternative splicing of serotonin 2C receptor reduces food intake. *EMBO Mol Med* 8: 878-894.
54. Compan V, Laurent L, Jean A, Macary C, Bockaert J, et al. (2012) Serotonin signaling in eating disorders. In: *WIREs Membrane Transport and Signaling* 715-729.
55. Bailer UF, Frank GK, Henry SE, Price JC, Meltzer CC, et al. (2007) Exaggerated 5-HT_{1A} but normal 5-HT_{2A} receptor activity in individuals ill with anorexia nervosa. *Biol Psychiatry* 10: 1090-1099.
56. Haahr M, Rasmussen P, Madsen K, Marner L, Ratner C, et al. (2012) Obesity is associated with high serotonin 4 receptor availability in the brain reward circuitry. *Neuroimage* 61: 884-888.
57. Ratner C, Ettrup A, Bueter M, Haahr ME, Compan V, et al. (2012) Cerebral markers of the serotonergic system in rat models of obesity and after Roux-en-Y gastric bypass. *Obesity (Silver Spring)* 20: 2133-2141.
58. Hardaway JA, Crowley NA, Bulik CM, Kash TL (2015b) Integrated circuits and molecular components for stress and feeding: Implications for eating disorders. *Gene, Brain and Behavior* 14: 85-97.
59. Steinglass J, Foerde K (2015) How does anorexia nervosa become resistant to change? In: *Managing severe and enduring anorexia nervosa*. London: Routledge.
60. Bonaventure P, Hall H, Gommeren W, Cras P, Langlois X, et al. (2000) Mapping of serotonin 5-HT₄ receptor mRNA and ligand binding sites in the post-mortem human brain. *Synapse* 36: 35-46.
61. Vilario MT, Cortes R, Mengod G (2005) Serotonin 5-HT₄ receptors and their mRNAs in rat and guinea pig brain: Distribution and effects of neurotoxic lesions. *J Comp Neurol* 484: 418-439.
62. Caille F, Morley TJ, Tavares AA, Papin C, Twardy NM, et al. (2013) Synthesis and biological evaluation of positron emission tomography radiotracers targeting serotonin 4 receptors in brain: [¹⁸F]MNI-698 and [¹⁸F]MNI -699. *Bioorg Med Chem Lett* 23: 6243-6247.
63. Marchetti-Gauthier E, Roman FS, Dumuis A, Bockaert J, Soumireu-Mourat B (1997) BIMU1 increases associative memory in rats by activating 5-HT₄ receptors. *Neuropharmacology* 36: 697-706.
64. Carlezon WA, Duman RS, Nestler EJ (2005) The many faces of CREB. *Trends Neurosci* 28: 436-445.
65. Koob GF, Nestler EJ (1997) The neurobiology of drug addiction. *J Neuropsychiatry Clin Neurosci* 9: 482-497.
66. Nestler EJ (2005) The neurobiology of cocaine addiction. *Sci Pract Perspect* 3: 4-10.
67. Chen JC, Chen PC, Chiang YC (2009) Molecular mechanisms of psychostimulant addiction. *Chang Gung Med J* 32: 148-154.
68. Nestler EJ (2004) Historical review: Molecular and cellular mechanisms of opiate and cocaine addiction. *Trends Pharmacol Sci* 25: 210-218.
69. Rocha BA, Searce-Levie K, Lucas JJ, Hiroi N, Castanon N, et al. (1998) Increased vulnerability to cocaine in mice lacking the serotonin-1B receptor. *Nature* 393: 175-178.
70. Couceyro PR, Evans C, McKinzie A, Mitchell D, Dube M, et al. (2005) Cocaine and amphetamine-regulated transcript (CART) peptides modulate the locomotor and motivational properties of psychostimulants. *J Pharmacol Exp Ther* 315: 1091-1100.
71. Rogge GA, Jones DC, Green T, Nestler E, Kuhar MJ (2009) Regulation of CART peptide expression by CREB in the rat nucleus accumbens in vivo. *Brain Res* 1251: 42-52.
72. Nestler EJ (2013) Treating the brain deep down: Brain surgery for anorexia nervosa? *Nat Med* 19: 678-679.
73. Amigo J, Diaz A, Pilar-Cuellar F, Vidal R, Martin A, et al. (2016) The absence of 5-HT₄ receptors modulates depression - and anxiety-like responses and

- influences the response of fluoxetine in olfactory bulbectomised mice: Adaptive changes in hippocampal neuroplasticity markers and 5-HT_{1A} autoreceptor. *Neuropharmacology* 111: 47-58.
74. Barrot M, Olivier JD, Perrotti LI, DiLeone RJ, Berton O, et al. (2002) CREB activity in the nucleus accumbens shell controls gating of behavioral responses to emotional stimuli. *Proc Natl Acad Sci U S A* 99: 11435-11440.
75. Steinglass JE, Figner B, Berkowitz S, Simpson HB, Weber EU, et al. (2012) Increased capacity to delay reward in anorexia nervosa. *J Int Neuropsychol Soc* 18: 773-780.
76. Brockmeyer T, Grosse Holtforth M, Bents H, Herzog W, Friederich HC (2013) Lower body weight is associated with less negative emotions in sad autobiographical memories of patients with anorexia nervosa. *Psychiatry Res*.
77. Wierenga C, Ely A, Bischoff-Grethe A, Bailer U, Simmons A, et al. (2014) Are extremes of consumption in eating disorders related to an altered balance between reward and inhibition? *Front Behav Neurosci* 8.
78. Gross C, Zhuang X, Stark K, Ramboz S, Oosting R, et al. (2002) Serotonin_{1A} receptor acts during development to establish normal anxiety-like behaviour in the adult. *Nature* 416: 396-400.
79. Conductier G, Dusticier N, Lucas G, Cote F, Debonnel G, et al. (2006) Adaptive changes in serotonin neurons of the raphe nuclei in 5-HT₄ receptor knock-out mouse. *Eur J Neurosci* 24: 1053-1062.
80. Vitousek K, Manke F (1994) Personality variables and disorders in anorexia nervosa and bulimia nervosa. *J Abnorm Psychol* 103: 137-147.
81. Kaye W, Strober M, Klump KL (2003) *Neurobiology of eating disorders*. New York: Oxford University Press.
82. Steinglass JE, Sysko R, Mayer L, Berner LA, Schebendach J, et al. (2010) Pre-meal anxiety and food intake in anorexia nervosa. *Appetite* 55: 214-218.
83. Amat J, Baratta MV, Paul E, Bland ST, Watkins LR, et al. (2005) Medial prefrontal cortex determines how stressor controllability affects behavior and dorsal raphe nucleus. *Nat Neurosci* 8: 365-371.
84. Robbins TW (2005) Controlling stress: How the brain protects itself from depression. *Nat Neurosci* 8: 261-262.
85. Euston DR, Gruber AJ, McNaughton BL (2012) The role of medial prefrontal cortex in memory and decision making. *Neuron* 76: 1057-1070.
86. Micali N, Solmi F, Horton NJ, Crosby RD, Eddy KT, et al. (2015) Adolescent eating disorders predict psychiatric, high-risk behaviors and weight outcomes in young adulthood. *J Am Acad Child Adolesc Psychiatry* 54: 652-659.
87. Jean A, Laurent L, Doly S, Ratner C, Dusticier N, et al. (2012b) Decision-making to eat after stress depends on the serotonin 4 and 1A receptors in the medial prefrontal cortex and dorsal raphe. In: 42ème Annual Meeting, Society for Neuroscience. New-Orleans, USA.
88. Walsh BT (2013) The enigmatic persistence of anorexia nervosa. *Am J Psychiatry* 170: 477-484.
89. Foerde K, Steinglass JE, Shohamy D, Walsh BT (2015) Neural mechanisms supporting maladaptive food choices in anorexia nervosa. *Nat Neurosci* 18: 1571-1573.
90. Solway A, Botvinick MM (2012) Goal-directed decision making as probabilistic inference: A computational framework and potential neural correlates. *Psychol Rev* 119: 120-154.
91. Wikler A (1961) On the nature of addiction and habituation. *Br J Addict Alcohol Other Drugs* 57: 73-79.
92. Hagen A, Manahan-Vaughan D (2017) The serotonergic 5-HT₄ receptor: A unique modulator of hippocampal synaptic information processing and cognition. *Neurobiol Learn Mem* 138: 145-153.
93. Segu L, Lecomte MJ, Wolff M, Santamaria J, Hen R, et al. (2010) Hyperfunction of muscarinic receptor maintains long-term memory in 5-HT₄ receptor knock-out mice. *PLoS ONE* 5: e9529.
94. Khodayar E, Oryan S, Nasehi M, Zarrindast MR (2016) Effect of nucleus accumbens shell 5-HT₄ receptors on the impairment of ACPA-induced emotional memory consolidation in male Wistar rats. *Behav Pharmacol* 27: 12-21.
95. Gershon MD (2016) 5-HT₄-mediated neuroprotection: a new therapeutic modality on the way? *Am J Physiol Gastrointest Liver Physiol* 310: G766-767.
96. Houghton LA, Heitkemper M, Crowell M, Emmanuel A, Halpert A, et al. (2016) Age, gender and women's health and the patient. *Gastroenterology*.
97. Briejer MR, Bosmans JP, Van Daele P, Jurzak M, et al. (2001) The in vitro pharmacological profile of prucalopride, a novel enterokinetic compound. *Eur J Pharmacol* 423: 71-83.

Citation: Compan V (2017) Serotonin 4 Receptors: A Cornerstone in Anorexia Nervosa? Autism Open Access 7: 207. doi:10.4172/2165-7890.1000207

OMICS International: Open Access Publication Benefits & Features

Unique features:

- Increased global visibility of articles through worldwide distribution and indexing
- Showcasing recent research output in a timely and updated manner
- Special issues on the current trends of scientific research

Special features:

- 700+ Open Access Journals
- 50,000+ editorial team
- Rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at major indexing services
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.omicsonline.org/submit/>